

ARBUZOV, A.Ye., akademik; SHISHKIN, V.Ye.

Interaction of imidoethers with alkyl halides. Dokl. AN SSSR 141
no.2:349-352 N '61.
(MIRA 14:11)

1. Kazanskiy khimiko-tehnologicheskiy institut im. S.M.Kirova.
(Ethers) (Halides)

ARBUZOV, A.Ye., akademik; SHISHKIN, V.Ye.

Rearrangement mechanism of imido ethers. Dokl. AN SSSR 141
no.3:611-612 N '61. (MIRA 14:11)

1. Kazanskiy khimiko-tehnologicheskiy institut im. S.M.
Kirova.

(Imides) (Ethers)

AREUZOV, A. YE., VALITOVA, E. G.

Naphtylenesalkyl esters of phosphorus acid.

Khimiya i Primeneniya Fosfororganicheskikh Soyedineniy (Chemistry and application of organophosphorus compounds) A. YE. AREUZOV, Ed.
Publ. by Kazan Affil. Acad. Sci. USSR, Moscow 1962, 632 pp.

Collection of complete papers presented at the 1959 Kazan Conference on
Chemistry of Organophosphorus Compounds.

ARBUZOV, A. YE.

PHASE I BOOK EXPLOITATION

SOV/6034

Konferentsiya po khimii i primeneniyu fosfororganicheskikh soyedineniy. 2d,
Kazan', 1959.

Khimiya i primeneniye fosfororganicheskikh soyedineniy; trudy (Chemistry
and Use of Organophosphorus Compounds; Conference Transactions) Moscow,
Izd-vo AN SSSR, 1962. 630 p. Errata slip inserted. 2800 copies printed.

Sponsoring Agency: Akademiya nauk SSSR. Kazanskiy filial.

Resp. Ed.: A. Ye. Arbuzov, Academician; Ed. of Publishing House: L. S.
Povarov; Tech. Ed.: S. G. Tikhomirova.

PURPOSE: This collection of conference transactions is intended for chemists,
process engineers, physiologists, pharmacists, physicians, veterinarians,
and agricultural scientists.

COVERAGE: The transactions include the full texts of most of the scientific
papers presented at the Second Conference on the Chemistry and Use of

Card 1/14

ARBUZOV, A.Ye.; VALITOVA, F.G.

Investigations in the field of ~~the~~ -diphenyl- β -picrylhydrazine.
Izv. AN SSSR Otd.khim.nauk no.2:354 F '62. (MIRA 15:2)

1. Khimicheskiy institut Kazanskogo filiala AN SSSR.
(Hydrazine)

ARBUZOV, A.Ye., akademik

N.N. Zinin, an outstanding Russian chemist; 150th anniversary of
his birth. Vest. AN SSSR 32 no.9:68-77 6 '62. (MIRA 15:9)
(Zinin, Nikolai Nikolaevich, 1812-1880)

ARBUZOV, A.Ye.

Great Russian scientist M.V.Lomonosov. Vop.ist.est.i tekhn.
no.12:3-21 '62. (MIRA 15:4)
(Lomonosov, Mikhail Vasil'evich, 1711-1765)

ARBUZOV, A.Ye., akademik; CHZHAN TSZIN-LIN [Chang Ching-ling]

Interaction of certain substituted triarylchloro- and bromo-methanes with salts of dialkylphosphorous acids. Dokl. AN SSSR 144 no.5:1039-1041 Je '62. (MIRA 15:6)

1. Kazanskiy filial Akademii nauk SSR.
(Methane) (Phosphorous acid)


S/020/62/147/001/015/022
B106/B101

AUTHORS: Arbuzov, A. Ye., Academician, Valitova, F. G., Il'yasov, A. V.,
Kosyrev, B. M., Yablokov, Yu. V.

TITLE: Study of the free radical α,α -diphenyl- β -pentaphenyl-cyclopentadienyl hydrazyl by the e.p.r. method


PERIODICAL: Akademiya nauk SSSR. Doklady, v. 147, no. 1, 1962, 99-102

TEXT: The e.p.r. spectrum of the free radical α,α -diphenyl- β -pentaphenyl-cyclopentadienyl hydrazyl (I) was studied both in solution and in its crystalline state. The synthesis of I was:



Card 1/4

Study of the free radical ...

S/020/62/147/001/015/022
B106/B101

(I). Data for the radical: yield 70-80%; ✓/✓

small bright-orange crystals with a melting point $> 180^{\circ}\text{C}$ (decomposition); soluble in benzene, chloroform, alcohol, acetonitrile, glacial acetic acid and dioxane. In dilute solutions ($< 10^{-3}$ moles/l), the spectra show a hyperfine structure, the analysis of which proves that the unpaired electron in I remains mainly on the nitrogen atoms. A comparison of the e.p.r. spectrum of I with the spectrum of the α,α -diphenyl- β -picryl hydrazyl radical (DPPH) showed that the additional hyperfine structure is due solely to the protons of the α -phenyl groups. It may be explained by the interaction of the unpaired electron with the 2,4,6-protons of one of the two α -phenyl groups. The value obtained for the constant a of hyperfine coupling was 1.7 oersteds, and for ΔH_n 1.1 oersted. The relative Card 2/4

S/020/62/147/001/015/022
B106/B101

Study of the free radical ...

stability of related free radicals from the e.p.r. spectra are estimated by the method of J. A. Weil, K. V. Sane, J. M. Kinkade (J. Phys. Chem., 65, 710 (1961)) showed that I is chemically more stable than DPPH. Its stability may be due to steric factors reducing the possibility of chemical reactions with other substances. The values obtained from the e.p.r. spectra of I in finely crystalline state, which may contain solvent, were 15.7 ± 0.3 oersteds for ΔH at 295°K , 10.5 ± 0.3 oersteds at 77°K , 1.43 for r at 295°K , and 1.45 at 77°C ($r = \langle \Delta H^4 \rangle^{1/4} / \langle \Delta H^2 \rangle^{1/2}$). The g-tensor at 295°K is: $g_1 = 2.0039 \pm 0.0001$, $g_2 = 2.0051 \pm 0.0001$, and $g_3 < g_1$. The considerable difference between these values and the g-factor of DPPH suggests that the molecular structure of the free radical considerably affects the residual spin - orbital coupling and anisotropy of the g-factor. There are 3 figures and 1 table. The most important English-language references are: M. M. Chen, K. V. Sane et al., J. Phys. Chem., 65, 713 (1961); B. Kubo, K. Tomita, J. Phys. Soc. Japan, 9, 888 (1954); F. K. Kneubuhl, J. Chem. Phys., 33, 1074 (1960).

Card 3/4

Study of the free radical ...

S/020/62/147/001/015/022
B106/B101

ASSOCIATION: Fiziko-tehnicheskiy institut Kazanskogo filiala Akademii nauk SSSR (Physicotechnical Institute of the Kazan' Branch of the Academy of Sciences USSR); Khimicheskiy institut im. A. Ye. Arbuzova Akademii nauk SSSR (Chemical Institute imeni A. Ye. Arbuzov of the Academy of Sciences USSR)

SUBMITTED: August 8, 1962

Card 4/4


S/020/62/147/004/017/027
B107/B186

AUTHORS: Arbuzov, A. Ye., Academician, Valitova, F. G.,
Il'yasov, A. V., Kozyrev, B. M., Yablokov, Yu. V.

TITLE: Electron paramagnetic resonance in solutions of some free
radicals of the phosphono-hydrazyl series

PERIODICAL: Akademiya nauk SSSR: Doklady, v. 147, no. 4, 1962, 839-842

TEXT: The resonance spectra of the following radicals were studied:




molar solutions in acetonitrile and chloroform were studied. A Р3 -1301 (RE-1301) radiofrequency spectrometer with a 9330 Mc frequency of the magnetic field was used. In all cases, a hyperfine structure of five equidistant lines was caused by interaction of the unpaired electron with the two N¹⁴ atoms. The spectrum is described by the spin Hamiltonian:

Card 1/3

S/020/62/147/004/017/027
B107/B186

Electron paramagnetic resonance ...


$\hat{\chi} = g\beta HS + A_1 \hat{S}I_{N_1} + A_2 \hat{S}I_{N_2}$, where β is the Bohr magneton, $g \approx g$
 $(\alpha, \alpha\text{-diphenyl-}\beta\text{-picryl hydrazyl}) = 2.0036$, H is the value of the static magnetic field, $S = 1/2$; $I_{N_1} = I_{N_2} = 1$. The constants A_1 and A_2 , and the width δH between maximum and minimum of the first derivative of the individual hyperfine structure line were obtained through comparison with theoretically plotted curves, using the given parameters. Calculated data agreed well with those obtained by experiments. $A_1 + A_2$ values found for phosphono-hydrazyls (maximum: 11.4 oe in azetonitrile, minimum: 9.4 in chloroform) were considerably less than the known value of 17.52 oe established for $\alpha, \alpha\text{-diphenyl-}\beta\text{-picryl-hydrazyl}$. A hyperfine structure caused by the P^{31} nucleus was not found. The production of phosphono-hydrazyls followed the reaction $(C_6H_5)_2^H N - N - P(C_6H_5)_2 \xleftarrow{H} (C_6H_5)_2^H N - NH_2 \xrightarrow{R_2P(O)Cl} (C_6H_5)_2^H N - N - PR_2$; and oxidation:



Card 2/3

S/020/62/147/004/017/027
B107/B186

Electron paramagnetic resonance ...



physical properties of phosphono-hydrazyls were tabulated. There are 1 figure and 2 tables.

ASSOCIATION: Khimicheskiy institut im. A. Ye. Arbuzova Akademii nauk SSSR (Chemical Institute imeni A. Ye. Arbuzov of the Academy of Sciences USSR); Fiziko-tehnicheskiy institut Kazanskogo filiala Akademii nauk SSSR (Physicotechnical Institute of the Kazan' Branch of the Academy of Sciences)

SUBMITTED: September 15, 1962

Card 3/3

ARBUZOV, A.Ye., akademik

Address at an out-of-town session of the Department of Chemical Sciences of the Academy of Sciences of the U.S.S.R. by Academician A.E.Arbusov. Izv.Kazan.fil. AN SSSR. Ser.khim.nauk no.6:3-10 '61. (MIRA 16:5)
(Phosphorus-organic compounds)

ARBUZOV, Aleksandr Yerminingel'dovich

[A.M.Butlerov, a great Russian chemist; the centenary of
the theory of chemical composition] A.M.Butlerov, velikii
russkii khimik; k 100-letiiu teorii khimicheskogo stro-
eniia. Moskva, Izd-vo AN SSSR, 1961. 42 p. (MIRA 16:8)
(Butlerov, Aleksandr Mikhailovich, 1828-1886)

ARBUZOV, A.Ye.; CHZHAN TSZIN-LIN [Chang Ching-ling]

Interaction of some substituted triaryl halomethanes with dialkyl phosphites. Report No.1: Interaction of p-alkoxy-substituted triphenylhalomethanes with dialkyl phosphites, Izv. AN SSSR. Ser. khim. no.11:1934-1941 N '63.

Interaction of some substituted triaryl halomethanes with dialkyl phosphites. Report No.2: Interaction of halo-substituted triphenyl halomethanes with dialkyl phosphites. Ibid.:1941-1944

Interaction of some substituted triaryl halomethanes with dialkyl phosphites. Report No.3: Interaction of crystalline triphenyl-chloromethane with sodium diethyl phosphite. Ibid.:1945-1946

1. Kazanskiy khimiko-tehnologicheskiy institut imeni S.M. Kirova.
(MIRA 17:1)

"APPROVED FOR RELEASE: 06/05/2000

CIA-RDP86-00513R000101920003-5

KRTAYEV, Yu.P.; TROYEPOL'SKAYA, T.V.; ARBUZOV, A. Ye.

Syntheses of heterocyclic compounds based on E. Fisher's reaction. Part 3: Catalysts of an "abnormal" course of reaction. Zhur. ob. Khim. 34 no.6:1835-1843 Je '64. (MIRA 17:7)

APPROVED FOR RELEASE: 06/05/2000

CIA-RDP86-00513R000101920003-5"

ARBOUZOV, A. Ye.; SHISHKIN, V. Ye.

Alcoholysis of imidoester haloalkylates. Zhur. ob. khim. 34 no.11:
3579-3582 N '64 (MIRA 18:1)

1. Kazanskiy khimiko-tehnologicheskiy institut imeni S.M. Kirova.

ARBUZOV, A.Ye.; SHAPSHINSKAYA, O.M.

Reactions of exchange decomposition of metallic derivatives of acid amides. Report No.2: Interaction of sodium and silver salts of benzamide with monochloromethyl, monobromomethyl, and monochloromethylethyl ethers. Trudy KKHTI no.30:22-27 '62.

(MIRA 16:10)

LAVROV, M.I.; NUZHIN, M.T., prof., otv.red.; MARKOV, M.V., prof., red.; DUBYAGO, A.D., prof., red.; ARBUZOV, A.Ye., akademik, red.; NORDEN, A.P., prof., red.; PIS'REV, V.I., prof., red.; TIKHVINSKAYA, Ye.I., prof., red.; FARYSHNIKOV, V.G., dotsent red.; KOLESNIKOVA, Ye. A., dotsent, red.; KOLOBOV, N.V., starshiy prepodavatel', red.; MOROZOV, D.G., dotsent, red.;

[Some statistical regularities of variable stars and their physical interpretation]. Nekotorye statisticheskie zakonomernosti u zatmennykh peremennnykh zvezd i ikh fizicheskoe istolkovar'ye. Kazan', 1955. 63 p. (Kazan. Universitet. Astronomicheskaiia observatoriia. Biulleten', no. 31) (MIRA 15:10)

1. Rektor Kazanskogo ordena Trudovogo Krasnogo Znameni gosudarstvennogo universiteta im. V.I.Ulyanova-Lenina (for Nuzhin).
2. Prorektor po nauchnoy rabote Kazanskogo ordena Trudovogo Krasnogo Znameni gosudarstvennogo universiteta im. V.I.Ulyanova-Lenina (for Markov).

KUZNETSOV, Vladimir Ivanovich; ARBUZOV, A.Ye., akademik, otv. red.;
KATRENKO, D.A., red.

[Advances in the field of catalytic organic synthesis]
Razvitiye kataliticheskogo organicheskogo sinteza. Mo-
skva, Nauka, 1964. 433 p. (MIRA 17:12)

DANILOV, S.N., glav. red.: ARBUZOV, A.Ye., red.; VVEDENSKIY, A.A.,
red.; VENUS-DANILOVA, E.D., red.; ZAKHAROVA, A.I., red.;
IOFFE, I.S., red.; KÁVERZNEVA, Ye.D., red.; LUTSENKO, I.F.,
red.; MISHCHENKO, K.P., red.; NEMTSOV, M.S., red.; PETROV,
A.A., red.; FREYDLINA, R.Kh., red.; SHEMYAKIN, M.M., red.;
SHUKAREV, S.A., red.; YUR'YEV, Yu.K., red.

[Biologically active compounds] Biologicheski aktivnye
soedineniya. Moskva, Nauka, 1965. 305 p.

(MIRA 18:7)


DANILOV, S.N., *glav. red.*; ZAKHAROVA, A.I., *red.*; ARBUZOV, A.Ye.,
red.; VVEDENSKIY, A.A., *red.*; VENUS-DANILova, E.D., *red.*;
IOFFE, I.S., *red.*; KAVERZNEVA, Ye.D., *red.*; LUTSENKO,
I.F., *red.*; MISHCHENKO, K.P., *red.*; NEMTSEV, M.S., *red.*;
PETROV, A.A., *red.*; FREYDLINA, R.Kh., *red.*; SHEMYAKIN,
M.M., *red.*; SHCHUKAREV, S.A., *red.*; YUR'YEV, Yu.K., *red.*


[Problems of organic synthesis] Problemy organicheskogo
sinteza. Moskva, Nauka, 1965. 323 p. (MIRA 18:8)


ARBUZOV, A.Ye.; SHISHKIN, V.Ye.; TYULENEV, S.S.


Imido ethers. Part 1: First haloalkylates of imido thio ethers.
Zhur. org. khim. 1 no.8:1442-1444 Ag '65. (MIRA 18:11)


1. Kazanskiy khimiko-tehnologicheskiy institut imeni Kirova.














	1ST AND 2ND ORDERS	3RD AND 4TH ORDERS
CA	PROCESSES AND PROPERTIES INDEX	
Materials Index	Chemical Index	
Open	Close	

10

Oxide of α - Δ^3 -carene. B. A. ARBUZOV AND B. M. MIKHAILOV. *J. Russ. Phys.-Chem. Soc.* 62, 807-16 (1930).— α - Δ^3 -Carene (I) belongs to the group of bicyclic terpenes, widely distributed in nature, which was discovered by Simonsen (*C. A.* 16, 2017). Δ^3 -Carene and Δ^2 -carene, which was found by S. in the essential oil of *Andropogon Juncuana* (*C. A.* 17, 81), are the first natural terpenes having the carene nucleus:

$\text{Me}_2\text{C}(\text{CH}_2)\text{CH}_2\text{CH}_2\text{CMe}_2\text{CH}_2\text{CH}_2$ (I) and $\text{Me}_2\text{C}(\text{CH}_2)\text{CH}_2\text{CH}_2\text{CMe}_2\text{CH}_2\text{CH}_2$.

S. (l. c.) oxidized I with KMnO_4 in alk. soln. to careneglycol, m. 60-70°, [α]_D 16.05°, and with HgCl_2 in AcOH to a glycol m. 80-1°, optically inactive in alc. and CHCl_3 (Pillay and S., *C. A.* 22, 1080). This isomerism of glycols was presumed by him to be either a *cis-trans* or optical isomerism, wherefore the glycol m. 60-70° was named α -carene- α -glycol, and the glycol m. 80-1° α -carene- β -glycol. By the action of dil. H_2SO_4 the β -glycol loses 1 mol. of H_2O with formation of a mixt. of *p*-cymene and


Epoxy α - Δ^3 -carene (II), which has an odor resembling limonene


and proved to be a very stable compd. giving no reaction with Br in CHCl_3 , and forming no hydrate on shaking for several days with dil. H_2SO_4 . The uncommon stability of II as contrasted with the high reactivity of the oxides of α -pinene and norpinene seems to be little understood, and is here proposed to be investigated. The Δ^2 -carene isolated

OVER

ASA-11A METALLURGICAL LITERATURE CLASSIFICATION									
ECONOMIC INFORMATION									
SEARCHED	INDEXED	SERIALIZED	FILED	SEARCHED					
SEARCHED	INDEXED	SERIALIZED	FILED	SEARCHED	INDEXED	SERIALIZED	FILED	SEARCHED	INDEXED
S	L	W	A	T	R	M	K	N	O

from turpentine gives on oxidation with $\text{Br}_2\text{O}_2\text{H}$ (Prilezhaev, C. A. 24, 607) 70% of the oxide of d,Δ^1 -carene (III), which differs sharply from II prepd. by S., i.e., II (fraction 1) b.p. 150-37°, d₄₀²⁰ 0.901, II (fraction 2) b.p. 165-9°, d₄₀²⁰ 0.9704, [α]_D²⁰ -30.10; III b.p. 70-81°, b.p. 120-7°, d₄₀²⁰ 0.9101, [α]_D²⁰ 13.05, m./w 1.124; III is a light liquid with a strong blue oil odor, and gives readily with dil. H_2SO_4 a hydrate, the main fraction of which is cryst. The d -carene α -glycol of S. m. 80-90°, optically inactive in CHCl_3 . The formation of β -glycol by hydration of III indicates for the α - and β -glycols of d,Δ^1 carene *cis-trans*, and not optical, isomerism, whereby the lower melting α -glycol is the *cis*, while the β -glycol is the *trans*-isomer (Nemtkin, C. A. 20, 2820; Verkade, et al., C. A. 23, 2055). The formation of identical glycols by hydration of an oxide and by oxidation of the corresponding unsatd. hydrocarbon with H_2O_2 in AcOH , as in the case of I, was previously observed (Sword, C. A. 19, 2042); Prilezhaev, C. A. 4, 910; Steerwein, C. A. 20, 2074). *Exptl. part.* -- From 4200 g. of Russian turpentine of unknown compn. was obtained by distn. in vacuo 815 g. (20.1%) of carene fractions, from which was isolated the pure d,Δ^1 -carene, b.p. 57-8°, b.p. 171-2.5°, d₄₀²⁰ 0.9248, m. 14.30°, n_D²⁰ 1.55, m./w 1.157. The filtrate of I was prepd. by known methods (C. A. 23, 4057). After recryst. from a mixt. of CHCl_3 and MeOH it decomps. 145°. Δ^1 -Carene- α -glycol was prepd. by methods developed by Simonsen (loc. cit.), Semmler and Schiller (C. A. 22, 241), and Krestinskii and Solodikii (C. A. 23, 4534), m. 70-1°. Δ^1 -Carene- β -glycol was prepd. according to Simonsen (C. A. 22, 1000), m. 80.5-90.5°. The oxide of I was obtained by pouring with stirring an ice-cold soln. of 87 g. of I in 300 cc. of anhyd. Et_2O into an ice-cold soln. of $\text{Br}_2\text{O}_2\text{H}$ in 1200 cc. of Et_2O contg. 10.3 g. of active O_2 ; after 20 hrs. of standing the reaction mass was washed 2 times with alkali, dried with fused K_2CO_3 , the Et_2O expelled and the residue (90 g.) fractionated at 12.5 mm. pressure, which finally produced the fraction b.p. 70-80°, or b.p. 120-7°, yield 21.7 g. or 70%. For the hydration of the oxide of Δ^1 -carene, 5 g. of the latter were shaken 1 hr. at room temp. with 25 cc. of 1% H_2SO_4 , the ppt. of the glycol was filtered off, washed with H_2O and dried in a vacuum desiccator over H_2SO_4 ; recryst. it m. 80-90°; yield 50%. The work is being continued. CHAS. BLANC








1ST AND 2ND ORDERS	PROCESSES AND PROPERTIES INDEX	3RD AND 4TH ORDERS																																																				
<p><i>Ca</i></p> <p>Presence of conjugated bonds in abietic acid. B. ABRUZOV. <i>J. Gen. Chem. (U. S. S. R.)</i> 2, 800-13 (1932).—While the presence of 2 double bonds in abietic acid (I) was conclusively demonstrated, their position was not made clear. The 2 double bonds show different reactivity giving easily the dihydro and difficultly the tetrahydro deriv. of I, and showing unlike speed in the oxidation with $\text{Br}_2\text{O}_2\text{H}$. From the formula of I proposed by Kuzicka (R. and Pfeiffer, <i>C. A.</i> 20, 421; R. et al., <i>C. A.</i> 25, 3657) there could be assumed the presence of conjugated bonds, an assumption here investigated by the action of maleic anhydride (II) on I. I, m. 166°, $[\alpha]_D -78.9^\circ$ (alc.), obtained from rosin acids by isomerization with HCl, was heated 4 hrs. at 170° in a sealed tube with II in dry C_6H_6, extd. with Et_2O and crystld. by adding petr. ether to incipient turbidity, giving an addn. compnd., $\text{C}_{17}\text{H}_{18}\text{O}_3$ (III), crystals from 100% AcOH, m. 227°. III, treated with NaOH and then acidified, gave the corresponding acid, which on heating was reconverted without melting to the anhydride, m. 227°. III (4 g.) with 6 g. powd. Se, heated 24 hrs. at 280-320°, then extd. with $\text{Br}_2\text{O}_2\text{H}$ and distd. in vacuo, produced 80% retene, crystals from MeCN, m. 97-8°, which with CrO_3 in AcOH gave retenequinone, m. 100-7°. The formation of retene does not indicate the position of the double bonds in I, and could be thus explained: (1) at the temp. of dehydrogenation (300-20°) III is decompd. into II and I, which is then converted to retene; (2) the cleavage of the new cycle takes place, because of its position in the mol. of III; (3) the newly formed ring generally can not remain during the dehydrogenation. The 1st case is excluded, because III heated 30 min. at 300-20° in a metal bath split off no II, the reaction beginning at 385-400° with the formation of $(\text{CH}_3\text{CO})_2\text{O}$, m. 118.5-9.5°. The other 2 cases are closely related and will require addnl. material for their interpretation. III (1.5 g.), treated with the calcd. amt. of aq. NaOH and then with 0.8 g. KMnO_4 in 120 cc. H_2O, extd. with CO_2, evapd., scidifed, filtered, washed and dried over H_2SO_4 produced 1.2 g. $\text{C}_{17}\text{H}_{18}\text{O}_5$ crystals from H_2O, m. 191-2°, which is presumed to be the monolactone, and not the anhydride, of the expected dihydroxytricarboxylic acid. It was interesting to investigate whether the</p>																																																						
<p style="text-align: center;">ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th colspan="2" style="text-align: left;">ECONOMIC CLASSIFICATION</th> <th colspan="2" style="text-align: right;">TECHNICAL SUBJECT</th> </tr> <tr> <th colspan="2"></th> <th style="text-align: center;">INDUSTRIAL</th> <th style="text-align: center;">SCIENTIFIC</th> </tr> <tr> <th>CLASS</th> <th>SECTION</th> <th>INDUSTRIAL</th> <th>SCIENTIFIC</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>0</td> <td>0</td> <td>0</td> </tr> <tr> <td>1</td> <td>1</td> <td>1</td> <td>1</td> </tr> <tr> <td>2</td> <td>2</td> <td>2</td> <td>2</td> </tr> <tr> <td>3</td> <td>3</td> <td>3</td> <td>3</td> </tr> <tr> <td>4</td> <td>4</td> <td>4</td> <td>4</td> </tr> <tr> <td>5</td> <td>5</td> <td>5</td> <td>5</td> </tr> <tr> <td>6</td> <td>6</td> <td>6</td> <td>6</td> </tr> <tr> <td>7</td> <td>7</td> <td>7</td> <td>7</td> </tr> <tr> <td>8</td> <td>8</td> <td>8</td> <td>8</td> </tr> <tr> <td>9</td> <td>9</td> <td>9</td> <td>9</td> </tr> </tbody> </table>			ECONOMIC CLASSIFICATION		TECHNICAL SUBJECT				INDUSTRIAL	SCIENTIFIC	CLASS	SECTION	INDUSTRIAL	SCIENTIFIC	0	0	0	0	1	1	1	1	2	2	2	2	3	3	3	3	4	4	4	4	5	5	5	5	6	6	6	6	7	7	7	7	8	8	8	8	9	9	9	9
ECONOMIC CLASSIFICATION		TECHNICAL SUBJECT																																																				
		INDUSTRIAL	SCIENTIFIC																																																			
CLASS	SECTION	INDUSTRIAL	SCIENTIFIC																																																			
0	0	0	0																																																			
1	1	1	1																																																			
2	2	2	2																																																			
3	3	3	3																																																			
4	4	4	4																																																			
5	5	5	5																																																			
6	6	6	6																																																			
7	7	7	7																																																			
8	8	8	8																																																			
9	9	9	9																																																			


ability of I to give III depends on the method of prepn. of I, because some authors presume that the acids obtained from resin do not contain double bonds, but decomp. with the rupture of the ring and the formation of double bonds in the process of production by the action of HCl or vacuum distn. (Tyutyunnikov and Perstnev, C. A. 26, 3395). I, obtained by vacuum distn. of various rosins and the primary resinous acids heated with II, produced in 5 expts. III; m. 227°, identical with III described above, while I, m. 173-4°, $[\alpha]_D -19.81^\circ$, obtained from French rosin by vacuum distn., and I, m. 158-61°, $[\alpha]_D = 0^\circ$, from the rosin of *Pinus sibirica* heated 24 hrs. at 270° with II, gave no addn. products. The exptl. data conclusively prove the presence of conjugated double bonds in I. The work is being continued.

CHAR. BLANC










"APPROVED FOR RELEASE: 06/05/2000


CIA-RDP86-00513R000101920003-5




APPROVED FOR RELEASE: 06/05/2000

CIA-RDP86-00513R000101920003-5"

1ST AND 2ND ORDERS												3RD AND 4TH ORDERS														
PROCESSES AND PROPERTIES INDEX												1G														
CONVENTIONAL LITERATURE OPEN MATERIALS INDEX	CA	<p>The structure of abietic acid, II. The synthesis of 1,3-dimethylcyclohexane-1,2,3-tricarboxylic acid. B. A. Arbusov and O. M. Staphanikaya. <i>Zhur. Nauk. Tekhn. Chern.</i>, No. 3, 19-34 (1935); <i>Ber.</i> 66B, 437-42 (1935); cf. <i>C. A.</i> 27, 2088. —By oxidation of abietic acid (I) with KMnO₄ Ruzicka <i>et al.</i> (<i>C. A.</i> 26, 431) obtained an acid (II), C₁₁H₁₆O₆, m. 218-9° (tri-Me ester, m. 75°), presumably 1,6-dimethylcyclohexane-1,2,3-tricarboxylic acid (<i>C. A.</i> 25, 3657). The work of Vocke (<i>C. A.</i> 26, 4350) and Haworth, <i>et al.</i> (<i>C. A.</i> 26, 4337) indicated the structure 1,3-dimethylcyclohexane-1,2,3-tricarboxylic acid. Ruzicka later (<i>C. A.</i> 27, 2134) suggested a formula for I based on the latter structure of II. A. and S. now report an attempt to synthesize II. Condensation of trimethylene bromide with 2 mols. of NaCMe(C₂H₅)₂ followed by hydrolysis gave α,α'-dimethylphthalic acid (III), a mixt. of the para and anti forms. As by-products in the condensation were formed a colorless liquid of unknown compn., b.p. 100-2, n_D²⁰ 1.4335, d₄²⁰ 1.0067, C₁₁H₁₆O₄, and a colorless liquid, C₁₁H₁₆O₆, b.p. 137-8°, n_D²⁰ 1.4360, d₄²⁰ 1.0040, mol. refraction 67.07, probably di-Et methylketohydroxypropanoic ester (IV). IV hydrolyzed to an acid, b.p. 175-8°, not obtained pure. III was converted to di-Et α,α'-dimethyl-α,α'-dibromoformate (V), colorless liquid, b.p. 182.5-4°, n_D²⁰ 1.4888, d₄²⁰ 1.4284, mol. refraction 81.19. V with NaC(C₂H₅)₂ gave tetra-Et 1,3-dimethylcyclohexane-1,2,3-tetracarboxylate (VI), b.p. (lit.) 8°, d₄²⁰ 1.1054, n_D²⁰ 1.4613, mol. refraction 90.26. With VI was also produced di-Et 1,3-dimethylpenta-1,4-diene-1,5-dicarboxylate (VII), b.p. 138.9°, n_D²⁰ 1.4775, d₄²⁰ 1.0451, mol. refraction 64.04. Hydrolysis of VII gave 2 stereoisomeric 1,5-dimethylpenta-1,4-diene-1,5-dicarboxylic acids, m. 165.8° and m. 161-3°. VI on hydrolysis gave the corresponding acid which rapidly lost CO₂. Elimination of CO₂ in vacuo at 100-5° gave an acid which could not be crystl. and hence identification with II was not possible. Partition through the Pb and Hg salts did not succeed. The Ag salt reacted with MeI to give a tri-Me ester which boiled similarly to that from II but could not be crystl. The analyses of the tri-Me ester and the Ag salt corresponded to II.</p> <p style="text-align: right;">Lewis W. Butt</p>																								
			450-514 METALLURGICAL LITERATURE CLASSIFICATION												EXTRACTIVE METALS											
			GENERAL SUBJECT												GENERAL SUBJECT											
			1000001 HIT ONLY SEE												100001 HIT ONLY SEE											
			1000001 HIT ONLY SEE												100001 HIT ONLY SEE											





IV. $\text{CMe}_2\text{CH-CH}_2\text{Br}$ and $\text{CHMe}^{\prime}\text{CMe-CHO}$ in Et_2O are added to Mg, when an alcohol, $\text{C}_{10}\text{H}_{14}$, b.p. $82^{\circ}-83^{\circ}$, is obtained. This is dehydrated by KHSO_4 at 180° to yield an unidentified terpene, b.p. 160° , not identical with the expected (I), and condensing with maleic anhydride to afford a product, m.p. $70^{\circ}-71^{\circ}$. (I) obtained from pinene is shown to be identical with natural (I), and with that obtained synthetically by the method of Fischer and Löwenberg (A., 1933, 302). R. T.

R. T.


APPROVED FOR RELEASE: 06/05/2000

CIA-RDP86-00513R000101920003-5"


ARBOUZOW, B.A.


"Sur l'isomerisation de l'-pinene en terpene aliphatique (alloocimene). Communication V". Arbouzow, B.A. (p. 292)

SO: Journal of General Chemistry. (Zhurnal Obshchey Khimii) 1936, Vol. 6, No. 2




100 AMP 1IN 09103	PROCESSES AND PROPERTIES INDEX	100 AMP 6IN 09103																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																
<p><i>BC</i> <i>a-3</i></p> <p>Structure of abietic acid. II. Attempted synthesis of 1:3-dimethylcyclohexane-1:2:3-tricarboxylic acid. B. A. ANAN'YEV and O. N. SOKARSKINSKAYA. <i>J. Gen. Chem. Russ.</i> 1936, 6, 404-410). —CH₂Br-CH₂-CH₂Br and CNaMe(CO₂Et)₂ in EtOH (at the b.p.) yield the Et₂ ester of α-dimethylpentane-α,ω-dicarboxylic acid (I), together with by-products, amongst which OEt-CH₂-CH₂-CH₂CMe(CO₂Et)₂ is identified. (I) and aq. EtOH-KOH (5 hr. at the b.p.) afford α-dimethyl-pimelic acid, which, with red P and Br (2 hr. at 100°), followed by EtOH (3-5 hr. at the b.p.), yields Et₂α-dibromo-α-dimethylpimelate, b.p. 183.5—184°/2-4 c.n., and this is condensed with CNa₂(CO₂Et)₂ in EtOH (100°; 3 hr.) to give the Et₂ ester of 1:3-dimethylcyclohexane-1:2:3-tetracarboxylic acid (II), b.p. 206—208°/2 mm., together with some Et₂α-dimethyl-Δ^5-pentadiene-α,ω-dicarboxylate, b.p. 138—139°/2 mm., from which the acid CH₂(CHCMe(CO₂H))₂, m.p. 168—171°, is obtained by hydrolysis. The acid obtained by hydrolysis of (II) (EtOH-KOH) yields a glassy mass, C₁₁H₁₆O₄ (Me₂ ester, b.p. 168—169°/0.7 mm.), when heated in vac. at 160—165°. The structure of the final product was not definitely established. R. T.</p>																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		
<p>ASBILIA METALLURGICAL LITERATURE CLASSIFICATION</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th colspan="10">1930-1934</th> <th colspan="10">1935-1939</th> </tr> <tr> <th colspan="5">1930-1934</th> <th colspan="5">1935-1939</th> <th colspan="5">1930-1934</th> <th colspan="5">1935-1939</th> </tr> <tr> <th>1</th><th>2</th><th>3</th><th>4</th><th>5</th> <th>6</th><th>7</th><th>8</th><th>9</th><th>10</th> <th>11</th><th>12</th><th>13</th><th>14</th><th>15</th><th>16</th><th>17</th><th>18</th><th>19</th><th>20</th><th>21</th><th>22</th><th>23</th><th>24</th><th>25</th><th>26</th><th>27</th><th>28</th><th>29</th><th>30</th><th>31</th><th>32</th><th>33</th><th>34</th><th>35</th><th>36</th><th>37</th><th>38</th><th>39</th><th>40</th><th>41</th><th>42</th><th>43</th><th>44</th><th>45</th><th>46</th><th>47</th><th>48</th><th>49</th><th>50</th><th>51</th><th>52</th><th>53</th><th>54</th><th>55</th><th>56</th><th>57</th><th>58</th><th>59</th><th>60</th><th>61</th><th>62</th><th>63</th><th>64</th><th>65</th><th>66</th><th>67</th><th>68</th><th>69</th><th>70</th><th>71</th><th>72</th><th>73</th><th>74</th><th>75</th><th>76</th><th>77</th><th>78</th><th>79</th><th>80</th><th>81</th><th>82</th><th>83</th><th>84</th><th>85</th><th>86</th><th>87</th><th>88</th><th>89</th><th>90</th><th>91</th><th>92</th><th>93</th><th>94</th><th>95</th><th>96</th><th>97</th><th>98</th><th>99</th><th>100</th><th>101</th><th>102</th><th>103</th><th>104</th><th>105</th><th>106</th><th>107</th><th>108</th><th>109</th><th>110</th><th>111</th><th>112</th><th>113</th><th>114</th><th>115</th><th>116</th><th>117</th><th>118</th><th>119</th><th>120</th><th>121</th><th>122</th><th>123</th><th>124</th><th>125</th><th>126</th><th>127</th><th>128</th><th>129</th><th>130</th><th>131</th><th>132</th><th>133</th><th>134</th><th>135</th><th>136</th><th>137</th><th>138</th><th>139</th><th>140</th><th>141</th><th>142</th><th>143</th><th>144</th><th>145</th><th>146</th><th>147</th><th>148</th><th>149</th><th>150</th><th>151</th><th>152</th><th>153</th><th>154</th><th>155</th><th>156</th><th>157</th><th>158</th><th>159</th><th>160</th><th>161</th><th>162</th><th>163</th><th>164</th><th>165</th><th>166</th><th>167</th><th>168</th><th>169</th><th>170</th><th>171</th><th>172</th><th>173</th><th>174</th><th>175</th><th>176</th><th>177</th><th>178</th><th>179</th><th>180</th><th>181</th><th>182</th><th>183</th><th>184</th><th>185</th><th>186</th><th>187</th><th>188</th><th>189</th><th>190</th><th>191</th><th>192</th><th>193</th><th>194</th><th>195</th><th>196</th><th>197</th><th>198</th><th>199</th><th>200</th><th>201</th><th>202</th><th>203</th><th>204</th><th>205</th><th>206</th><th>207</th><th>208</th><th>209</th><th>210</th><th>211</th><th>212</th><th>213</th><th>214</th><th>215</th><th>216</th><th>217</th><th>218</th><th>219</th><th>220</th><th>221</th><th>222</th><th>223</th><th>224</th><th>225</th><th>226</th><th>227</th><th>228</th><th>229</th><th>230</th><th>231</th><th>232</th><th>233</th><th>234</th><th>235</th><th>236</th><th>237</th><th>238</th><th>239</th><th>240</th><th>241</th><th>242</th><th>243</th><th>244</th><th>245</th><th>246</th><th>247</th><th>248</th><th>249</th><th>250</th><th>251</th><th>252</th><th>253</th><th>254</th><th>255</th><th>256</th><th>257</th><th>258</th><th>259</th><th>260</th><th>261</th><th>262</th><th>263</th><th>264</th><th>265</th><th>266</th><th>267</th><th>268</th><th>269</th><th>270</th><th>271</th><th>272</th><th>273</th><th>274</th><th>275</th><th>276</th><th>277</th><th>278</th><th>279</th><th>280</th><th>281</th><th>282</th><th>283</th><th>284</th><th>285</th><th>286</th><th>287</th><th>288</th><th>289</th><th>290</th><th>291</th><th>292</th><th>293</th><th>294</th><th>295</th><th>296</th><th>297</th><th>298</th><th>299</th><th>300</th><th>301</th><th>302</th><th>303</th><th>304</th><th>305</th><th>306</th><th>307</th><th>308</th><th>309</th><th>310</th><th>311</th><th>312</th><th>313</th><th>314</th><th>315</th><th>316</th><th>317</th><th>318</th><th>319</th><th>320</th><th>321</th><th>322</th><th>323</th><th>324</th><th>325</th><th>326</th><th>327</th><th>328</th><th>329</th><th>330</th><th>331</th><th>332</th><th>333</th><th>334</th><th>335</th><th>336</th><th>337</th><th>338</th><th>339</th><th>340</th><th>341</th><th>342</th><th>343</th><th>344</th><th>345</th><th>346</th><th>347</th><th>348</th><th>349</th><th>350</th><th>351</th><th>352</th><th>353</th><th>354</th><th>355</th><th>356</th><th>357</th><th>358</th><th>359</th><th>360</th><th>361</th><th>362</th><th>363</th><th>364</th><th>365</th><th>366</th><th>367</th><th>368</th><th>369</th><th>370</th><th>371</th><th>372</th><th>373</th><th>374</th><th>375</th><th>376</th><th>377</th><th>378</th><th>379</th><th>380</th><th>381</th><th>382</th><th>383</th><th>384</th><th>385</th><th>386</th><th>387</th><th>388</th><th>389</th><th>390</th><th>391</th><th>392</th><th>393</th><th>394</th><th>395</th><th>396</th><th>397</th><th>398</th><th>399</th><th>400</th><th>401</th><th>402</th><th>403</th><th>404</th><th>405</th><th>406</th><th>407</th><th>408</th><th>409</th><th>410</th><th>411</th><th>412</th><th>413</th><th>414</th><th>415</th><th>416</th><th>417</th><th>418</th><th>419</th><th>420</th><th>421</th><th>422</th><th>423</th><th>424</th><th>425</th><th>426</th><th>427</th><th>428</th><th>429</th><th>430</th><th>431</th><th>432</th><th>433</th><th>434</th><th>435</th><th>436</th><th>437</th><th>438</th><th>439</th><th>440</th><th>441</th><th>442</th><th>443</th><th>444</th><th>445</th><th>446</th><th>447</th><th>448</th><th>449</th><th>450</th><th>451</th><th>452</th><th>453</th><th>454</th><th>455</th><th>456</th><th>457</th><th>458</th><th>459</th><th>460</th><th>461</th><th>462</th><th>463</th><th>464</th><th>465</th><th>466</th><th>467</th><th>468</th><th>469</th><th>470</th><th>471</th><th>472</th><th>473</th><th>474</th><th>475</th><th>476</th><th>477</th><th>478</th><th>479</th><th>480</th><th>481</th><th>482</th><th>483</th><th>484</th><th>485</th><th>486</th><th>487</th><th>488</th><th>489</th><th>490</th><th>491</th><th>492</th><th>493</th><th>494</th><th>495</th><th>496</th><th>497</th><th>498</th><th>499</th><th>500</th><th>501</th><th>502</th><th>503</th><th>504</th><th>505</th><th>506</th><th>507</th><th>508</th><th>509</th><th>510</th><th>511</th><th>512</th><th>513</th><th>514</th><th>515</th><th>516</th><th>517</th><th>518</th><th>519</th><th>520</th><th>521</th><th>522</th><th>523</th><th>524</th><th>525</th><th>526</th><th>527</th><th>528</th><th>529</th><th>530</th><th>531</th><th>532</th><th>533</th><th>534</th><th>535</th><th>536</th><th>537</th><th>538</th><th>539</th><th>540</th><th>541</th><th>542</th><th>543</th><th>544</th><th>545</th><th>546</th><th>547</th><th>548</th><th>549</th><th>550</th><th>551</th><th>552</th><th>553</th><th>554</th><th>555</th><th>556</th><th>557</th><th>558</th><th>559</th><th>560</th><th>561</th><th>562</th><th>563</th><th>564</th><th>565</th><th>566</th><th>567</th><th>568</th><th>569</th><th>570</th><th>571</th><th>572</th><th>573</th><th>574</th><th>575</th><th>576</th><th>577</th><th>578</th><th>579</th><th>580</th><th>581</th><th>582</th><th>583</th><th>584</th><th>585</th><th>586</th><th>587</th><th>588</th><th>589</th><th>590</th><th>591</th><th>592</th><th>593</th><th>594</th><th>595</th><th>596</th><th>597</th><th>598</th><th>599</th><th>600</th><th>601</th><th>602</th><th>603</th><th>604</th><th>605</th><th>606</th><th>607</th><th>608</th><th>609</th><th>610</th><th>611</th><th>612</th><th>613</th><th>614</th><th>615</th><th>616</th><th>617</th><th>618</th><th>619</th><th>620</th><th>621</th><th>622</th><th>623</th><th>624</th><th>625</th><th>626</th><th>627</th><th>628</th><th>629</th><th>630</th><th>631</th><th>632</th><th>633</th><th>634</th><th>635</th><th>636</th><th>637</th><th>638</th><th>639</th><th>640</th><th>641</th><th>642</th><th>643</th><th>644</th><th>645</th><th>646</th><th>647</th><th>648</th><th>649</th><th>650</th><th>651</th><th>652</th><th>653</th><th>654</th><th>655</th><th>656</th><th>657</th><th>658</th><th>659</th><th>660</th><th>661</th><th>662</th><th>663</th><th>664</th><th>665</th><th>666</th><th>667</th><th>668</th><th>669</th><th>670</th><th>671</th><th>672</th><th>673</th><th>674</th><th>675</th><th>676</th><th>677</th><th>678</th><th>679</th><th>680</th><th>681</th><th>682</th><th>683</th><th>684</th><th>685</th><th>686</th><th>687</th><th>688</th><th>689</th><th>690</th><th>691</th><th>692</th><th>693</th><th>694</th><th>695</th><th>696</th><th>697</th><th>698</th><th>699</th><th>700</th><th>701</th><th>702</th><th>703</th><th>704</th><th>705</th><th>706</th><th>707</th><th>708</th><th>709</th><th>710</th><th>711</th><th>712</th><th>713</th><th>714</th><th>715</th><th>716</th><th>717</th><th>718</th><th>719</th><th>720</th><th>721</th><th>722</th><th>723</th><th>724</th><th>725</th><th>726</th><th>727</th><th>728</th><th>729</th><th>730</th><th>731</th><th>732</th><th>733</th><th>734</th><th>735</th><th>736</th><th>737</th><th>738</th><th>739</th><th>740</th><th>741</th><th>742</th><th>743</th><th>744</th><th>745</th><th>746</th><th>747</th><th>748</th><th>749</th><th>750</th><th>751</th><th>752</th><th>753</th><th>754</th><th>755</th><th>756</th><th>757</th><th>758</th><th>759</th><th>760</th><th>761</th><th>762</th><th>763</th><th>764</th><th>765</th><th>766</th><th>767</th><th>768</th><th>769</th><th>770</th><th>771</th><th>772</th><th>773</th><th>774</th><th>775</th><th>776</th><th>777</th><th>778</th><th>779</th><th>780</th><th>781</th><th>782</th><th>783</th><th>784</th><th>785</th><th>786</th><th>787</th><th>788</th><th>789</th><th>790</th><th>791</th><th>792</th><th>793</th><th>794</th><th>795</th><th>796</th><th>797</th><th>798</th><th>799</th><th>800</th><th>801</th><th>802</th><th>803</th><th>804</th><th>805</th><th>806</th><th>807</th><th>808</th><th>809</th><th>810</th><th>811</th><th>812</th><th>813</th><th>814</th><th>815</th><th>816</th><th>817</th><th>818</th><th>819</th><th>820</th><th>821</th><th>822</th><th>823</th><th>824</th><th>825</th><th>826</th><th>827</th><th>828</th><th>829</th><th>830</th><th>831</th><th>832</th><th>833</th><th>834</th><th>835</th><th>836</th><th>837</th><th>838</th><th>839</th><th>840</th><th>841</th><th>842</th><th>843</th><th>844</th><th>845</th><th>846</th><th>847</th><th>848</th><th>849</th><th>850</th><th>851</th><th>852</th><th>853</th><th>854</th><th>855</th><th>856</th><th>857</th><th>858</th><th>859</th><th>860</th><th>861</th><th>862</th><th>863</th><th>864</th><th>865</th><th>866</th><th>867</th><th>868</th><th>869</th><th>870</th><th>871</th><th>872</th><th>873</th><th>874</th><th>875</th><th>876</th><th>877</th><th>878</th><th>879</th><th>880</th><th>881</th><th>882</th><th>883</th><th>884</th><th>885</th><th>886</th><th>887</th><th>888</th><th>889</th><th>890</th><th>891</th><th>892</th><th>893</th><th>894</th><th>895</th><th>896</th><th>897</th><th>898</th><th>899</th><th>900</th><th>901</th><th>902</th><th>903</th><th>904</th><th>905</th><th>906</th><th>907</th><th>908</th><th>909</th><th>910</th><th>911</th><th>912</th><th>913</th><th>914</th><th>915</th><th>916</th><th>917</th><th>918</th><th>919</th><th>920</th><th>921</th><th>922</th><th>923</th><th>924</th><th>925</th><th>926</th><th>927</th><th>928</th><th>929</th><th>930</th><th>931</th><th>932</th><th>933</th><th>934</th><th>935</th><th>936</th><th>937</th><th>938</th><th>939</th><th>940</th><th>941</th><th>942</th><th>943</th><th>944</th><th>945</th><th>946</th><th>947</th><th>948</th><th>949</th><th>950</th><th>951</th><th>952</th><th>953</th><th>954</th><th>955</th><th>956</th><th>957</th><th>958</th><th>959</th><th>960</th><th>961</th><th>962</th><th>963</th><th>964</th><th>965</th><th>966</th><th>967</th><th>968</th><th>969</th><th>970</th><th>971</th><th>972</th><th>973</th><th>974</th><th>975</th><th>976</th><th>977</th><th>978</th><th>979</th><th>980</th><th>981</th><th>982</th><th>983</th><th>984</th><th>985</th><th>986</th><th>987</th><th>988</th><th>989</th><th>990</th><th>991</th><th>992</th><th>993</th><th>994</th><th>995</th><th>996</th><th>997</th><th>998</th><th>999</th><th>1000</th></tr> </thead></table>			1930-1934										1935-1939										1930-1934					1935-1939					1930-1934					1935-1939					1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700	701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800	801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	890	891	892	893	894	895	896	897	898	899	900	901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	990	991	992	993	994	995	996	997	998	999	1000
1930-1934										1935-1939																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																								
1930-1934					1935-1939					1930-1934					1935-1939																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																			
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700	701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800	801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	890	891	892	893	894	895	896	897	898	899	900	901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	990	991	992	993	994	995	996	997	998	999	1000																																											





(1) AND (2) ORDER		(3) AND (4) ORDER																															
PROCESSES AND PROPERTIES INDEX																																	
<p><i>B. ARBUZOV</i></p> <p>Some products of the dione synthesis with piperylene and benzene. N. Arbusov, Z. Zhur. org. i. khim., 1, 2278 (1937).—The hexadiene fraction of the bivinyl still residue from synthetic rubber mainly condenses with acrolein to give 2,5-dimethyl-3-cyclohexen-1-al (II) and with crotonaldehyde to give 2,5,6-trimethyl-3-cyclohexen-1-al (III). The piperylene fraction and acrolein give 5-methyl-3-cyclohexen-1-al (III). I condenses with MeEtCO to form a ketone which is probably RCH:CHCOEt, but may be RCH:CMcCOMe (R = 2,5-dimethyl-3-cyclohexen-1-yl), bp 132-4°, d₄²⁰ 0.9376, n_D²⁰ 1.4931, M. R. 60.1A. With Me₂CO, I gives an analogous compound, which is either RCH:CHCOEt or RCH:CH(COMe), bp 134-5°, d₄²⁰ 0.9143, n_D²⁰ 1.4921, M. R. 60.30. I and methyl oxide give RCH:CHCOCH₂COMe, bp 138-40°, d₄²⁰ 0.9121, n_D²⁰ 1.4900, M. R. 70.00. III and MeEtCO give R'CH:CHCOEt or R'CH:CMcCOMe, bp 140°, d₄²⁰ 0.9225, n_D²⁰ 1.4983, M. R. 65.02 (R' = 5-methyl-3-cyclohexen-1-yl). I and EtCHO give RCH:CHCH₂CHO, bp 120-1°, d₄²⁰ 0.9050, n_D²⁰ 1.4976, M. R. 60.00. III and EtCHO give R'CH:CHCH₂CHO, bp 110-12°, d₄²⁰ 0.9446, n_D²⁰ 1.4910, M. R. 49.42, and some 5-methyl-3-ethylalcohol. II condenses with EtCOAc and Na to give R'CH:CHCOEt, bp 124-5°, d₄²⁰ 0.9628, n_D²⁰ 1.4820, M. R. 67.44. The Et acetal of I, bp 117-19°, d₄²⁰ 0.9200, n_D²⁰ 1.4570, M. R. 62.85; the Bu acetal, bp 153-5°, d₄²⁰ 0.9003, n_D²⁰ 1.4011, M. R. 81.57. The Et acetal of III, bp 91-3°, d₄²⁰ 0.9280, n_D²⁰ 1.4544, M. R. 57.05. By the Grignard reaction with I the following alcs. are prep'd.: RCH(OH)Et (IV), bp 120-3°, d₄²⁰ 0.9140, n_D²⁰ 1.4826, M. R. 51.29; RCH(OH)Pr, bp 120-2°, d₄²⁰ 0.9200, n_D²⁰ 1.4830, M. R. 60.01; iso-BuCH(OH)R, bp 124-0°, d₄²⁰ 0.9378, n_D²⁰ 1.4840, M. R. 50.87; iso-C₁₁H₂₃OEt, bp 107-8°, d₄²⁰ 0.9380, n_D²⁰ 1.4842, M. R. 60.71. Oxidation of IV with K₂Cr₂O₇ gives the corresponding ketone, bp 100-2°, d₄²⁰ 0.9272, n_D²⁰ 1.4700; semicarbazone, m. 100°. II gives by the Grignard reaction R'CH(OH)Et (R' = 2,5,6-trimethyl-3-cyclohexen-1-yl), bp 116-8°, d₄²⁰ 0.9240, n_D²⁰ 1.4820, M. R. 56.16; R'CH(OH)Pr, bp 118-20°, d₄²⁰ 0.9112, n_D²⁰ 1.4822, M. R. 61.35 and iso-BuCH(OH)R', bp 132-4°, d₄²⁰ 0.9142, n_D²⁰ 1.4841, M. R. 65.76. All the compds. except the aldehydes derived from condensation with EtCHO have pleasant odors.</p> <p>H. M. Leicester</p>																																	
ASB-11-A METALLURGICAL LITERATURE CLASSIFICATION																																	
<table border="1"> <thead> <tr> <th colspan="2">STANDARD SUBJECT</th> <th colspan="2">SECONDARY SUBJECTS</th> <th colspan="2">EDITIONS</th> <th colspan="2">COUNTRY OF PUBLICATION</th> <th colspan="2">PUBLISHER OR SOURCE</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> <td>6</td> <td>7</td> <td>8</td> <td>9</td> <td>10</td> </tr> <tr> <td>11</td> <td>12</td> <td>13</td> <td>14</td> <td>15</td> <td>16</td> <td>17</td> <td>18</td> <td>19</td> <td>20</td> </tr> </tbody> </table>				STANDARD SUBJECT		SECONDARY SUBJECTS		EDITIONS		COUNTRY OF PUBLICATION		PUBLISHER OR SOURCE		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
STANDARD SUBJECT		SECONDARY SUBJECTS		EDITIONS		COUNTRY OF PUBLICATION		PUBLISHER OR SOURCE																									
1	2	3	4	5	6	7	8	9	10																								
11	12	13	14	15	16	17	18	19	20																								



Isomerization of terpenic oxides. II. Isomerization of α -pinene oxide in the Grignard reaction. B. A. Arbuzov, J. Gen. Chem. (U. S. S. R.) 9, 249-54 (1939); cf. C. A. 29, 8887. — The results of an independent study on the isomerization and reduction of condensation of α -pinene oxide (I) with the Grignard reagent (cf. C. A. 30, 2948) agree with the findings of Ritter and Russel (C. A. 30, 2948) that the product of reaction is an alkyl- or aryl-substituted camphorol and not $\text{CH}_2\text{CH}(\text{OH})\text{CH}(\text{OH})\text{CH}_2$.

$\text{CH}_2\text{CHCHMe}_2$ (II) as was shown by P. and V. The reaction of 40 g. Et_2Zn and 50 g. I-I at 100° and subsequent hydrolysis with H_2O and acid, in HCl proceeds without the isomerization and rupture of the pinane nucleus to give 18 g. of an *alc.*, CuH_2O , bp. 81.5-2.5°, d_4^{20} 0.9248, n_D^{20} 1.475, M. R. n_D 65.42, [α]_D 10.04 (cf. Cook, et al., C. A. 30, 2542). I with PhLi failed to react. III. Isomerization of the oxides of camphene, camphane and α^1 -carene in the Reformatsky reaction. *Ibid.* 248-71. — Studies on the isomerization of α -pinene oxide with the intermediate formation of campholenaldehyde (I) in the Reformatsky reaction are extended to addnl. bicyclic terpenes. Camphene oxide (III), nopinene oxide (III) and α^1 -carene oxide (IV) were prep'd. from the corresponding terpenes by oxidation with $\text{Ac}_2\text{O}/\text{Fe}^{+2}$ in Et_2O . Refluxing 25 g. II, m. 80-81°, [α]_D 10.02, in 50 ml. of dry CaH_2 with 0.2-0.3 g. ZnBr_2 , removal of the CaH_2 and redistn. of the residue afforded 20 g. I. II gave as sole product with Zn and

BuCH_2COEt (VI) an *ester*, CuH_2O_2 (VII), bp. 131.5-2.5°, d_4^{20} 1.0483, n_D^{20} 1.483, M. R. n_D 65.32, isopnd. to a solid. Zn and V on I, it appears to be the *E*-ester of β -hydroxy- δ -(2,2-dimethylbicyclo[2.2.1]hept-5-en-3)-propionic acid. III, bp. 68.5-9.5°, d_4^{20} 0.9083, n_D^{20} 1.478, M. R. n_D 44.36, does not react with ZnCl_2 and ZnBr_2 in Et_2O , but in CaH_2 it is isomerized into *dihydromyrcenol*, bp. 81-2°, d_4^{20} 0.9033, n_D^{20} 1.4812, M. R. n_D 44.96; *semicarbazone*, m. 152-3°, III with Zn and V gave a solid. *ester* (VII), CuH_2O_2 , bp. 137.8°, d_4^{20} 1.0278, n_D^{20} 1.48, M. R. n_D 60.31. It appears to be the *E*-ester of β -hydroxy- δ -(7,7-dimethylbicyclo[1.1.1]hept-5-en-2)-propionic acid. IV, bp. 80-81.5°, d_4^{20} 0.9578, n_D^{20} 1.460, M. R. n_D 44.34, [α]_D 11.05. The complex reaction of IV

Lub. Org. Chem.-, Chem. Sci. Res. Inst.
im. Butakov

ASR-5A METALLURGICAL LITERATURE CLASSIFICATION

ARBUZOV, B.A.; ABRAMOV, V.S.; DEVIATOV, YA. B.

"The Products of Condensation of Cyclones with p-Benzoquinone and Napthoquinone",
Zhur. Obshch. Khim., 9, No. 17, 1939. Laboratory of Organic Chemistry, Kazan'
Chemico-Technological Institute imeni S. M. Kirov. Rec'd 22 Feb 1939.

[redacted] Report U-1614, 3 Jan 1952.

"APPROVED FOR RELEASE: 06/05/2000


CIA-RDP86-00513R000101920003-5


ARBUZOV, B. A.

"Uber Die Struktur Der -Pimarsaure," (About the structure of Primary Acids), Iz. Ak.
Nauk USSR, Ot. Khim. Nauk, No. 1, 1940


APPROVED FOR RELEASE: 06/05/2000

CIA-RDP86-00513R000101920003-5"





Action of naphthazarin on hexadiene and piperylene.
B. A. Arbuzov and K. Nikonorov. *J. Gen. Chem.* (U. S. S. R.) 10, 649-52 (1940).—Condensation of hexadiene (I) and piperylene (II), resp., with naphthazarin (III) was carried out first in abs. alc. with the formation of resinous. When PhNO_2 was used instead of alc., and the mixt. was heated in a sealed tube at 100-70° for 2 hrs. *1,4-dimethyl-5,8-dihydroxyanthraquinone* was obtained from I and III, orange-red needles, m. 220-7°, in 73.7% yield, and *1-methyl-5,8-dihydroxyanthraquinone* from II and III, when heated in a sealed tube at 125-30° for 20 hrs., red needles, m. 220-7°, yield 42.7% (based on III).



When benzene was used as reaction medium, negative results were obtained. Condensation of III with alkoxime (IV) in abs. alc. in CO_2 atm. in a sealed tube at 80-90° gave a reaction product $\text{C}_{11}\text{H}_{12}\text{O}_2$, red needles, m. 157°, yield 20.5%. The same compd. was obtained when benzene was used, whereas no cryst. product could be isolated in the case of PhNO_2 . Though the structure for this compd. has not yet been proved, formula V is sug-

gested because of results obtained in previous studies on the condensation of α -naphthoquinone with IV.

10

ARBUZOV, B. A.

"On the Structure of *L*-Pimamic Acid," Dok. AN, 30, No. 8, 1941. State Univ. Kazan.
Butlerov Chem. Research Inst.

"APPROVED FOR RELEASE: 06/05/2000

CIA-RDP86-00513R000101920003-5

ARBUZOV, B. A., Butlerov, A. M.

"Structure of 1-pimamic Acid," Organic Chem Lab, Inst of Chem Research Investigation,
Kazan State Univ., Uchenye Zapiski Kazan Gosudarst Univ, Vol. 101, No. 1, 1941

W-257, 2 Mar 48 441699

APPROVED FOR RELEASE: 06/05/2000

CIA-RDP86-00513R000101920003-5"

ARBUSOW, B. A.


"The action of dihydronaphthalenes on cyclones." Arbusow, B. A., and Akhmed-Zade, J. A. (p. 211)

SO: Journal of General Chemistry (Zhurnal Obshchei Khimii) 1942, Vol 12, No 3-4.

ARBUSOW, B. A.

"Piperylcyclone and its derivatives." Arbusow, B. A., and Akhmed-Zade, J. A. (p. 218)

SO: Journal of General Chemistry (Zhurnal Obshchei Khimii) 1942, Vol 12, No 3-4.



ARBUZOV, Boris Aleksandrovich

"Use of Parachors and Dipole Moments for Studying the Fine Structure of Some
Organic Derivatives of Phosphorous, Boron, Nitrogen, and Silicon." Zhur. Obsch.
Khim., 13, Nos. 1-2, 1943

1ST AND 2ND ORDERS												3RD AND 4TH ORDERS											
PROCESS AND PROPERTIES INDEX																							
<p><i>C</i></p> <p>Addition products of dienes to toluene. B. A. Arbuzov and R. V. Kurnetsov. <i>Comp. rend. acad. sci. U. R. S. S.</i> 30, 311-13 (1943) (in English).—Toluene (I) and butadiene (II) were caused to react in the presence of finely divided Na and the products were cyclized and then dehydrogenated with S. Reaction of 300 g. I and 200 cc. II in presence of Na dust at 90° for 10 hrs. yielded 142 g. of a product (b_{10} 80-220°), whose fractionation yielded 1-phenyl-3-pentene (III), b_{10} 92-4°, n_D^{20} 1.5690, d_4^{20} 0.870; 3-phenyl-7,7-dimethyloctadiene (IV), b_{10} 140-2°, n_D^{20} 1.6108, d_4^{20} 0.880; PhCH₂(C₆H₅)₂, b_{10} 188-01°, n_D^{20} 1.5135, d_4^{20} 0.888; PhCH₂(C₆H₅)H, b_{10} 210-90°, n_D^{20} 1.5138, d_4^{20} 0.920. Cyclization of III yielded 95% of 1,2,3,4-tetrahydro-1-methylnaphthalene (V), which with S yielded 75% of dehydrogenation product, b_{10} 145°, n_D^{20} 1.6130, d_4^{20} 1.001. Similarly IV produced 90% of cyclization product, b_{10} 162-4°, n_D^{20} 1.5471, d_4^{20} 0.980, thought to be 1-<i>phenylhexahydronaphthalene</i>, whose dehydrogenation yielded no pos. result. Condensation of V with II at 100° with Na catalyst yielded 15% of a product (b_{10} 130-2°, n_D^{20} 1.5378, d_4^{20} 0.983) formulated as 1-(2-butenyl)-1,2,3,4-tetrahydronaphthalene, which was cyclized to a product b_{10} 148-50°, n_D^{20} 1.5548, d_4^{20} 1.0008, which resinated on attempted dehydrogenation with S. Conjugated dienes with substituents on the terminal C atoms reacted much less readily with I than did II. Thus 300 g. I and 250 g. 2,4-heptadiene with 20 g. Na at 70° for 10 hrs. yielded only 40 g. of a product from which was isolated 1-<i>phenyl-3-methyl-3-hexene</i>, b_{10} 100-10°, n_D^{20} 1.4905, d_4^{20} 0.873. J. W. Perry</p>																							
ASB-SLA METALLURICAL LITERATURE CLASSIFICATION																							
2ND REVISIOVA												3RD REVISIOVA											
SECOND REV		SECOND REV ORV GSC		MILLION		SECOND REV ORV GSC		MILLION		SECOND REV ORV GSC		MILLION											
SOURCE	NO.	S	P	D	S	P	D	S	P	D	S	P	D										
D	SL	SI	AV	10	11	D	P	D	P	D	S	P	D										

		PROCESSES AND PROPERTIES INDEX																		3RD AND 4TH COLUMNS	
		1ST AND 2ND ORDERS									3RD AND 4TH ORDERS										
COMPOUND ELEMENTS		<i>Cd</i>																		COMPOUND ELEMENTS	
		1ST AND 2ND ORDERS									3RD AND 4TH ORDERS										
MATERIALS INDEX		Use of dihalo ethers for the synthesis of oxygen-containing cyclic acids. I. Ali-Zade and B. A. Arshamov. <i>J. Gen. Chem. (U. S. S. R.)</i> 13, 113-23 (in English, 123-4) (1943).—Contrary to the data of Dox and Yoder (<i>C. A.</i> 36, 1862), the authors showed that the reaction of $(BrCH_2)_2O$ with metallic derivs. of malonic ester yields the normal product, $[(HO_2C)CHCH_2]_2O$. Mg (40 g.) and 100 cc. abs. EtOH were treated with 250 g. $CH_3(CO_2Et)_2$ (after addn. of $\frac{1}{4}$ of the malonate the mixt. was heated on a steam bath for 45 min. and 1 cc. CCl_4 added to initiate the reaction). After complete soln. of the Mg, the flask was cooled and 150 g. $(BrCH_2)_2O$ was added dropwise, after which the mixt. was heated on a steam bath for 3 hrs., cooled, treated with dil. H_2SO_4 , extd. with Et ₂ O, the latter dried and distd., yielding 100 g. of a heavy oil, b.p. 188-71°, d ₄ ²⁰ 1.1128, n _D ²⁰ 1.4382, and 20 g. of material b.p. 171-303°, which crystallized on standing and m. 80-2° (from aq. EtOH); the former compd. is bis(2,2-dicarbethoxyethyl) ether, while the 2nd substance was not investigated further. Hydrolysis of the ester by KOH in aq. EtOH gave a tetrabasic acid, m. 147-0°. The ester (20 g.), 2.7 g. Mg and 20 cc. EtOH were heated on a steam bath and treated with 0.6 cc. CCl_4 to initiate the reaction, after which the heating was continued until the Mg disappeared; on cooling the mixt. was treated with 10 g. iodine in EtOH and heated on a steam bath for 8 hrs.; after the usual treatment with acidified water and extn. with Et ₂ O, there was obtained 8 g. <i>tetra-Et tetrahydro-3,3,6,6-tetracarboxylate</i> , b.p. 171-4°, d ₄ ²⁰ 1.1293, n _D ²⁰ 1.4401. Similar reaction with CH_3Br gave <i>tetra-Et tetrahydro-3,3,5,5-tetracarboxylate</i> , b.p. 170-82°, d ₄ ²⁰ 1.1131, n _D ²⁰ 1.4100, while $(CH_3)_2Br$ gave <i>tetra-Et hexamethylene-3,3,6,6-tetracarboxylate</i> , b.p. 158-60°, d ₄ ²⁰ 1.1110, n _D ²⁰ 1.4388; $Cl(CH_3)_2Br$ gave $(EtO_2C)_2CHCH_2OCH_2C(ClO_2)CH_2CH_2Cl$, b.p. 102-6°, d ₄ ²⁰ 1.1026, n _D ²⁰ 1.4398; $(CH_2=CHCH_2)_2CO$ gave 35% of $OCH_2C(CO_2Et)_2CH_2CH_2CH_2CO$.																			
		MATERIALS INDEX		AB-SLA METALLURGICAL LITERATURE CLASSIFICATION																	
1ST AND 2ND ORDERS									3RD AND 4TH ORDERS												
SEARCHED		SEARCHED ONE ONLY ONE																		SEARCHED ONE ONLY ONE	
		SEARCHED									SEARCHED										
INDEXED		INDEXED																		INDEXED	
		INDEXED									INDEXED										
FILED		FILED																		FILED	
		FILED									FILED										

$(CO_2Et)_2CH_2$ (I), b, 101-4°, d₄₄ 1.1000, n_D 1.4619;
 $(BrCH_2)_2O$ gave 40% of $O.CH_2.C(CO_2Et)_2CH_2.O.CH_2.C-$

$(CO_2Et)_2CH_2$ (II), b, 188-9°, d₄₄ 1.1415, n_D 1.4630.


$(BrCH_2CH_2)_2O$ was prep'd. by treating a cooled mixt. of 12 g. dry pyridine and 44 g. $O(CH_2CH_2OH)$, with 88 g. PBr_3 ; after standing overnight the product was distd. in tomo., washed, dried and redistd., b, 110°. Na (2.3 g.) in 50 cc. EtOH was treated with 15 g. of *tetra-Bu*₄*N*₊*Br*₋*tetracarboxylate* in 26 cc. EtOH; after heating on a steam bath for 1 hr., the mixt. was treated with 8 g. $(BrCH_2CH_2)_2O$ and heated on a steam bath for 7 hrs., the EtOH distd. off, the residue decomposed with water and extd. with Et₂O; on diazn. of the ext. there was obtained 8 g. *tetra-Bu*₄*N*₊*Br*₋*octamethyltriacarboxylate-4,5,6,7-tetracarboxylate*, b, 208-12°, d₄₄ 1.1572, n_D 1.4661. $[CH_2CH(CO_2Et)_2]_2$ (from $(CH_2)_2Br$) and $CH_2(CO_2Et)_2$ in the form of its Mg salt was treated with $(BrCH_2CH_2)_2O$, yielding *tetra-Bu*₄*N*₊*Br*₋*octamethyltriacarboxylate-4,5,6,7-tetracarboxylate*, in 17% yield, b, 190-4°, m. 75-7°. The Na compd. from 19.3 g. Na, 300 cc. EtOH and 80 g. $CO(CH_2CO_2Et)_2$ was treated with 80 g. $(BrCH_2CH_2)_2O$, heated for 18 hrs., the NaBr sepd.; the soin. heated for 62 addnl. hrs. with several seprs. of NaBr; the EtOH was then distd., the residue treated with water and extd. with Et₂O, yielding 34 g., b, 188-63, d₄₄ 1.1562, n_D 1.4615, of an oil which corresponded to $CH_2CH_2O.CH_2CH_2O.C(CH_2CO_2Et)CCO_2Et$.

Hydrolysis of this substance with alc. KOH gave a monobasic acid, m. 87-9°, $C_6H_{10}O_4$ which appears to be either $CH_2CH_2O.CH_2CH_2O.C(CH_2CO_2H)CH_2$ or $CH_2CH_2O.CH_2CH_2O.CMe(CO_2H)CH_2$.

O. M. K.

ARBUZOV, B. A., Rafikov, S. R., and Zoroastrova, V. M.

"Preparation of the Dinitrile of Adipic Acid" Inst Org Chem, Acad Sci, Moscow,
Bul Acad Sci URSS, Class Sci Chim, 1945



"APPROVED FOR RELEASE: 06/05/2000


CIA-RDP86-00513R000101920003-5

ARBUZOV, B. A.

"On the Preparation of Dinitrile of Adipic Acid," Iz. Ak. Nauk SSSR, Otdel. Khim. Nauk, No. 2, 1945. Inst. of Org. Chem, Acad Sci, USSR.

APPROVED FOR RELEASE: 06/05/2000

CIA-RDP86-00513R000101920003-5"



ARBUZOV, B. A., Zelinskiy, N. D. and Shuikin, N. I.


"Oxidation of Cyclohexene by Selenious Acid and a New Method of Preparation of 1, 3-cyclohexadiene," Bul. Acad. Sci. URSS, Classe sci. chim. 1945, 163-6.


SO; Chemical Abstracts, Vol. 40, No. 12, 20 Jun 46

ARBUZOV, B. A.

The isotropyl ester of benzenephosphorous acid. [✓]
B. Arbuзов, G. Kh. Kamal, and O. N. Belorossova (Kazan
Chem. Tech. Inst.), *J. Gen. Chem. (U.S.S.R.)* 15, 766-9
(1945). - When iso-PrOH and PhNMe₂ in dry Et₂O are
cooled and treated with PhPCl₃ in a CO₂ atm., they form a
distillate, 60% of which is *di-iso-Pr* benzenephosphonite
(I), b.p. 121-2°, d₂₀²⁰ 1.0103, d₄²⁰ 0.9952, n_D²⁰ 1.5021, and
15% *iso-Pr* phenylisopropylphosphinate (II), b.p. 146-7°,
d₂₀²⁰ 1.0957, d₄²⁰ 1.0813, n_D²⁰ 1.4929, formed by isomerization
of I. When I is heated to 150° in the presence of iso-PrI
(III) it liberates MeCH₂CH₃ and forms *phenylisopropyl-*
phosphinic acid, m. 61-2°, which gives cryst. Na, K, Ca,
and Ba salts. When I and III are allowed to stand 10 days
at room temp., 11.1% isomerization to II occurs. Addn.
of a little PhNMe₂ increases the isomerization to 95%.

H. M. Leicester





ARBUZOV, B. A.

Reduction of nitriles of dibasic acids over Raney nickel.
B. A. Arbuзов and E. A. Пузултая. *Bull. acad. sci. U.R.S.S., Class. sci. chim.*, 1946, 65-70 (in Russian).—

The dinitriles of adipic, sebacic, and succinic acids are readily reduced over Raney Ni at 75-80° to the corresponding amino nitriles under 0.5-0.8 atm. H₂; adiponitrile is reduced to (C₁₁H₂)₂(NH₂)₂ at 100-10° and 80-100 atm. H₂. The catalyst was made from Al-Ni alloy by treatment with boiling 25% NaOH; after decanting, a fresh batch of alkali was used and the mist, heated 1 hr. at 90°; this was repeated 2-3 times and the catalyst was washed free of alkali by H₂O and abs. EtOH and was used immediately. Adiponitrile (10 g.), 150 g. BuOH, and 5 g. catalyst were treated with H₂ (0.5-0.8 atm.) at 75-80° 10 hrs.; after filtration and removal of the BuOH *in vacuo*, the residue was dried, with H₂O and 0.5 g. unreacted dinitrite was removed. Treatment of the residue with BzCl resulted in

Isolation of 93% Bz deriv. of the resulting *ε*-aminocapronitrile, m. 92-3° (from 70% alc.); hydrolysis by HCl gave *ε*-aminocaproic acid-HCl, which gave the free acid, m. 200-2°; the same yield was obtained if the catalyst was added in portions during the course of reduction. Succinonitrile (8 g.), 150 cc. BuOH, and 4 g. catalyst similarly treated with H₂ 19 hrs. at 75-80° gave the Bz deriv. of γ -aminobutyronitrile (no data given, but a general statement indicated a yield of 90%), as an oil, which on hydrolysis by HCl gave γ -aminobutyric acid, m. 181-2°; HCl salt, m. 134-5°; chloroplatinate, m. 218-20°. Sebaconitrile (10 g.), 150 g. BuOH, and 5 g. catalyst were hydrogenated as above for 26 hrs.; filtration, removal of the BuOH, and treatment of the residue with BzCl gave (yield not given) the Bz deriv. of *ε*-aminocaproic acid, m. 103-5°, which on hydrolysis by HCl gave the free acid, m. 185-7°; chloroplatinate, m. 208-302° (decompn.). Adiponitrile (50 g.), 450 g. BuOH, and 17 g. catalyst heated 2 hrs. to 110° at 90 atm. initial H₂ pressure (15 atm. final) gave 1.3 g. unreacted dinitrite, and 46 g. (85.7%) hexamethylenediamine, b. 200-4°; 3.5 g. of tar was formed. Increase of the temp. to 150° (14 hrs.) dropped the yield to 26.8%; halving of the catalyst amt. dropped the yield to 64%. The high-pressure runs were made in a rotating Bergius autoclave.

G. M. Kosolapoff

JARBUZOV, B. A.

Condensation products of dimethylol adipimide and trimethylol derivatives of imides of other dibasic acids with aromatic compounds. B. A. Arbusov and D. A. Livshits. *Zvest. Akad. Nauk S.S.R., Otdel. Khim. Nauk* 1946, 391-402.—Adipimide (25 g.), heated on a water bath to 70-80° with 42.5 g. 30% formalin, 25 ml. H₂O, and 2.5 g. K₂CO₃ until soln. took place, then an addnl. 2 hrs., and cooled, gave 71% *N,N*'-dimethyloladipimide, (CH₂CH₂CONHCH₂OH)₂, (I), m. 148-0.7° (from H₂O). To 8.5 g. C₆H₆ and 42 ml. concd. H₂SO₄ was added gradually 10 g. I at 8-10° and after 3 days at room temp. the mixt. was poured on ice, and the resulting solid washed with H₂O and alc., boiled with H₂O, and extd. with hot Et₂CO, giving an amorphous insol. portion, m. 225-7°, and a sol. portion, m. 174-5°, the total yield being 92%. The sol. product, C₂₁H₂₂O₂N₂, gives Ba(OH)₂ on oxidation with KMnO₄, and appears to be PhCH₂NHCO(CH₂)₂CONHCH₂Ph. The insol. product, C₁₁H₁₂O₂N₂, gives terephthalic acid on oxidation by KMnO₄, and appears to be either a polymer —(C₆H₅CH₂NHCO(CH₂)₂CONHCH₂)_n— or a cyclic product *p*-C₆H₄(CH₂NHCOCH₂CH₂)₂. AcNHPh (2.6 g.) in 20 ml. concd. H₂SO₄ (treated at 0° with 2 g. I, allowed to stand 2 days, poured on ice, and the resulting solid washed with water and recrystd. from AcOH), yielded 1.7 g. (41%) C₁₁H₁₂O₂N₂, m. 244-

1.5°, which on boiling with 20% HCl gave *p*-H₂NCH₂/CH₂NH₂Cl, decompt. above 300°; the product is apparently *p*-AcNH₂C₆H₄CH₂NHCO(CH₂)₂CONHCH₂CH₂NH₂/Ac, as on boiling with 30% NaOH it gives 88% *p*-H₂NCH₂CH₂NHCO(CH₂)₂CONHCH₂CH₂NH₂, m. 173-4° (from EtOH). Coupling of the diazotate of this diamine with PhNMe₂ in HCl gave the corresponding [*p*-Me₂NCH₂H₂N:NC₆H₄CH₂NHCO(CH₂)₂]_n, m. 160-70°; 2HCl, black-green, m. 185-6° (from AcOH); similar coupling with 2-naphthol in the presence of Na₂CO₃-NaOH¹ gave an analogous 2-hydroxy-1-naphthyl deriv., m. 244-6° (by pptn. from H₂SO₄ by H₂O). Condensation of *p*-acetoluidide with I in H₂SO₄, as above, gave 55% C₂₁H₂₂N₂O₂, m. 233-4° (from AcOH), which probably has a structure similar to that of the AcNHPh deriv., while *p*-acetoluidide gave a similar product, m. 245-6° (from EtOH). Condensation of I with 2-acetophenylhalide in H₂SO₄ at 12-15° gave a similar product, m. above 300°, insol. in the usual org. solvents. Addn. of 2 g. I to cold 2.7 g. *p*-O₂NCH₂OH in 20 ml. concd. H₂SO₄, followed by 24 hrs.' standing, gave 80% [5,2-(NO₂)(OH)C₆H₄CH₂NHCO(CH₂)₂]_n, m. 230-10°, yellow sol. in alkalies, AcOH, and EtOH; hydrolysis by concd. HCl gave 2-hydroxy-5-nitrobenzylamine-HCl, m. 240° (from H₂O). 2-Naphthol (2.8 g.) and 2 g. I in 20 ml. EtOH and 20 ml. H₂O treated with 30 drops of concd. H₂SO₄ gave after standing 3 days 50% colorless [2-HOC₆H₄CH₂NHCO(CH₂)₂]_n, m. 222-4° (from AcOH), which on hydrolysis,

(over)

		PROCESSES AND		PROPERTIES INDEX	
		1ST AND 2ND ORDERS			
<i>Ca</i>		<p>Allenic rearrangements. I. Isomerization of some methoxychloropentenes in the presence of metal chlorides. A. N. Pudovik and B. A. Arbuzov. <i>Izvst. Akad. Nauk S.S.R., Otdel Khim. Nauk</i> 1946, 227-37 (in Russian); cf. C.A. 42, 4073b.—The addn. of ClCH_2OMe to butadiene was improved over the procedure given by Strauss and Thiel (C.A. 31, 3031). Butadiene (20 g.) cooled to -15° was treated with 310 g. ClCH_2OMe, followed by 4-5 g. powd. ZnCl_2, the flask closed with a stopper provided with an outlet tube and pinch-clamp, and the mixt. carefully mixed by shaking and allowed to stand in an ice-bath 5-8 hrs., then 15-20 hrs. at room temp. When the pressure drops, the reaction is complete; the residual butadiene is removed and the residue, dild. with Et_2O, is washed with H_2O and distd. to give 135 g. 1-methoxy-3-chloro-4-pentene, bp 34-35°, $d_4^{20} 0.9090$, $n_D^{20} 1.4361$, 103 g. 1-methoxy-5-chloro-3-pentene, bp 38-39°, $d_4^{20} 1.0000$, $n_D^{20} 1.4540$, and 75 g. high-boiling products. The addn. does not go with the catalyst, even after 1 yr. standing. Neither the 3-(I), nor the 5-Cl isomer (II) shows isomerization on standing 1 yr. at room temp. in sealed tubes. When I and II were heated to 100° in sealed tubes, I isomerized to the extent of 16% in 27 hrs., while II gave but 3% isomerization, as detd. by n. If the temp. was raised to 140° the reaction was speeded up, but incipient decomp. limited the duration to 15 hrs., at which point I was 65.9% isomerized and II 7.9%. The isomerization was also studied in the presence of catalysts: 1% and 10% ZnCl_2, and 10% SnCl_4. ZnCl_2 (10%) enabled equil. to be reached in 24 hrs.; the mixt. then contained 80% II (equil. reached from both sides); when 1% ZnCl_2 is used the reaction is slower and</p> <p>even after 300 hrs. II is only 11.3% isomerized, while I is 63.1% isomerized. The results are given graphically. SnCl_4 (10%) caused a somewhat less extensive isomerization, but the reaction could not be followed accurately because of tar formation. Use of 10% AlCl_3 gave a greater heat evolution on mixing than was observed with the other catalysts, but its isomerization efficiency is poor: after 24 hrs. II is isomerized 11%, and 143%; 1% AlCl_3 is much less effective. If the addn. of butadiene to ClCH_2OMe is conducted using 1% AlCl_3 catalyst, the reaction mixt. gives a 30% total yield, composed of 23% II and 77% I. The problem of the possibility of primary addn. being at the 1,2-positions only cannot be solved from the available data: there is an indication that the 1,2-addn. is favored. The metal chloride-catalyzed isomerization probably proceeds similarly to the Friedel-Crafts reaction, involving formation of mesomeric ions of the methoxypentene and ZnCl_2-type ions, the isomerization being based on the common resonant ion. The lack of attainment of equil. when low percentages of the catalysts were used is explained by secondary reactions which give products incapable of isomerization. II. Action of sodium acetate on isomeric methoxychloropentenes, and hydrolysis of the resulting acetates. <i>Ibid.</i> 511-8.—The reaction of NaOAc with isomeric methoxychloropentenes is best interpreted on the customary basis of allylic reactions, with the secondary chloride reacting by the S_{N}^2 reaction mechanism, while the primary chloride reacts by both S_{N}^2 and to some extent by S_{E} mechanisms. 1-Methoxy-5-chloro-3-pentene (30 g.), 23 g. AcOH, and 25 g. dry NaOAc heated 2 hrs. at $120-30^\circ$ gave 2.3 g. 1-methoxy-4-penten-3-ol ac-</p>			
AS-A SLA METALLURGICAL LITERATURE CLASSIFICATION					
AS-AH STATUS		SUBJECT MAT. ONV. GRC		SECTION	
SEARCHED	INDEXED	SEARCHED	INDEXED	SEARCHED	INDEXED
SEARCHED	INDEXED	SEARCHED	INDEXED	SEARCHED	INDEXED

late (I), b₅ 63-5°, n_D²⁰ 1.4264, d₄²⁰ 0.9000, and 23.7 g. *1-methoxy-3-penten-5-ol acetate* (II), b₅ 82-5°, n_D²⁰ 1.4300, d₄²⁰ 0.9789. 1-Methoxy-3-chloro-4-pentene (30 g.), 23 g. AcOH, and 25 g. AcONa heated 2 hrs. at 120-30° gave 10.7 g. I and 16 g. II. II (10 g.), 10 g. AcOH, and 10 g. NaOAc heated 2 hrs. at 120-5° produced no change; the same result was obtained with I. I and II were likewise unaffected by boiling with Ag₂O 3 hrs. Hydrolysis of II by hot alc. KOH gave only *1-methoxy-3-penten-3-ol*, b₅ 97°, n_D²⁰ 1.4485, d₄²⁰ 0.9523; similar hydrolysis of I gave *1-methoxy-3-penten-3-ol*, b₅ 90-1°, n_D²⁰ 1.4373, d₄²⁰ 0.9300. Hydrolysis of the chlorides by alkali also goes without isomerization.

G. M. Kowalsoff

"APPROVED FOR RELEASE: 06/05/2000

CIA-RDP86-00513R000101920003-5

ARBUZOV, B. A. and Zoroastrova, V. M.


"The Reaction of N,N-dichlorobenzenesulfonamide and Butadiene in Ethanol". Compte Rend Acad Sci URSS, Vol. 5, 1946


"Simultaneous Action of Chlorine and Sodium Alcohols (As well as Chlorine and Acetic Anhydride) on Butadiene" Compte Rend Acad Sci URSS, VOL 53, 1946


APPROVED FOR RELEASE: 06/05/2000


CIA-RDP86-00513R000101920003-5"

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100																																																																								
	1ST AND 2ND GROUPS												3RD AND 4TH GROUPS																																																																																																																																																															
	PROCESSES AND PROPERTIES INDEX																																																																																																																																																																											
	10																																																																																																																																																																											
<p>100. The simultaneous action of chlorine and sodium al- cobolate (as well as chlorine and acetic anhydride) on butadiene. II. A. Arbusov and V. M. Zoroastrova. <i>Compt. rend. acad. sci. U.R.S.S.</i> 53, 41-4 (1946); cf. <i>C.A.</i> 39, 4834. —A detailed study of the addn. of Et hypohalite and mixed anhydrides of acetic and chloric acids to bivinyl in order to investigate the general phenomena of 1,2- and 1,4-addn. is reported. The following compds. were obtained by repeated fractionation of the products from an alc. soln. of Na with Cl and bivinyl until the resulting soln. was distinctly alk.: 1-chloro-2- ethoxy-3-butene (I), b. 130-4°, n_D²⁰ 1.4340, d₄²⁰ 0.9784, MR_D 35.80 (theoretical 35.00); 1-chloro-4-ethoxy-2- butene (II), b. 61-4°, n_D²⁰ 1.4480, d₄²⁰ 1.0070, MR_D 39.70; and (III) a product corresponding by analysis to a compd. formed by the addn. of 2 mols. of ROCl and 1 M bivinyl, b. 90-1°, n_D²⁰ 1.4530, d₄²⁰ 1.1300, MR_D 51.39 (theoretical, 51.27). For identification, I was transformed by alc. alkali into ethylcyprene, CH₃-C(OR)₂-CH=CH₂ (IV), b. 94-8°, n_D²⁰ 1.4400, d₄²⁰ 0.8300, and by the formation of MeCOCH₂CH₃ after the hydrolysis of IV. For identification II was converted by the action of alc. KOH into 1,4-diehtoxy-3-butene (V), b. 81°, d₄²⁰ 0.9001, n_D²⁰ 1.4208, MR_D 41.00 (theoretical 42.15). III is receiving further study to locate the positions of the EtO groups and the Cl atoms. It is certain from the above results, that the entering groups of EtOCl are added to both the 1,2- and 1,4-positions of butadiene. The simul- taneous action of Cl and Ac₂O in the presence of anhyd. NaOAc, on butadiene is likewise reported. Repeated fractions produced the following products: 1-chloro-3- acetoxy-3-butene (VI), b. 84-0°, b₄₀ 104.5-0.3°, n_D²⁰ 1.4530, d₄²⁰ 1.1300, MR_D 35.08 (theoretical, 35.00); 1-chloro-3-acetoxy-3-butene (VII), b. 82-4.5°, n_D²⁰ 1.4700, d₄²⁰ 1.1510, MR_D 30.00; a product (VIII) resulting from the addn. of 2 Cl atoms to VI or VII, b. 114-16°, n_D²⁰ 1.4700, d₄²⁰ 1.3010, MR_D 45.03 (theoretical 46.10); and a product (IX), m. 117.8°, resulting from the addition of 2 mols. AcOCl to bivinyl. VI was converted by solid KOH into 1,2-oxybivinyl (X), b. 65-7°, n_D²⁰ 1.4220, d₄²⁰ 0.8690. VII was not isolated in the pure state but was converted to 2-butene-1,4-diol (XI) and then by NaOAc into the diacetate, AcOCH₂CH=CHCH₂OAc (XII), m. 14-16°, b. 114°, d₄²⁰ 1.0842, n_D²⁰ 1.4470, MR_D 42.4 (theoretical, 41.98). VIII appears to be derived pre- dominately from VI. The structure of IX is likewise indefinite but appears to result from the simultaneous addn. of a mol. of AcOCl at the 1,2- and 1,4-positions. The Cl content of II, III, VI, VII, VIII, and IX confirms the former conclusions.</p> <p style="text-align: right;">R. R. Dunbar</p>																																																																																																																																																																												
<p>400-500 METALLURICAL LITERATURE CLASSIFICATION</p> <table border="1"> <thead> <tr> <th rowspan="2">ITEM NUMBER</th> <th colspan="12">GENERAL SUBJECT</th> <th colspan="12">SUBJECT</th> </tr> <tr> <th colspan="3">GENERAL SUBJECT</th> <th colspan="3">SUBJECT</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> <td>6</td> <td>7</td> <td>8</td> <td>9</td> <td>10</td> <td>11</td> <td>12</td> <td>13</td> <td>14</td> <td>15</td> <td>16</td> <td>17</td> <td>18</td> <td>19</td> <td>20</td> <td>21</td> <td>22</td> <td>23</td> <td>24</td> <td>25</td> <td>26</td> <td>27</td> <td>28</td> <td>29</td> <td>30</td> <td>31</td> <td>32</td> <td>33</td> <td>34</td> <td>35</td> <td>36</td> <td>37</td> <td>38</td> <td>39</td> <td>40</td> <td>41</td> <td>42</td> <td>43</td> <td>44</td> <td>45</td> <td>46</td> <td>47</td> <td>48</td> <td>49</td> <td>50</td> <td>51</td> <td>52</td> <td>53</td> <td>54</td> <td>55</td> <td>56</td> <td>57</td> <td>58</td> <td>59</td> <td>60</td> <td>61</td> <td>62</td> <td>63</td> <td>64</td> <td>65</td> <td>66</td> <td>67</td> <td>68</td> <td>69</td> <td>70</td> <td>71</td> <td>72</td> <td>73</td> <td>74</td> <td>75</td> <td>76</td> <td>77</td> <td>78</td> <td>79</td> <td>80</td> <td>81</td> <td>82</td> <td>83</td> <td>84</td> <td>85</td> <td>86</td> <td>87</td> <td>88</td> <td>89</td> <td>90</td> <td>91</td> <td>92</td> <td>93</td> <td>94</td> <td>95</td> <td>96</td> <td>97</td> <td>98</td> <td>99</td> <td>100</td> </tr> </tbody> </table>																								ITEM NUMBER	GENERAL SUBJECT												SUBJECT												GENERAL SUBJECT			1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100																					
ITEM NUMBER	GENERAL SUBJECT												SUBJECT																																																																																																																																																															
	GENERAL SUBJECT			SUBJECT			SUBJECT			SUBJECT			SUBJECT			SUBJECT			SUBJECT			SUBJECT																																																																																																																																																						
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100																																																																									









Parachors and structure of esters of phosphoric, phosphorous, and phosphonic acids. II. A. Arbutov and V. S. Vinogradova (A. M. Butlerov Research Inst., Kazan State Univ.). Bull. acad. sci. U.R.S.S., Classe sci. chim., 1947, 450-72 (in Russian); cf. C.A. 41, 20086.—(1) The following esters were synthesized and their surface tensions, γ (dyne/cm.), at 20° detd. by the method of max. bubble pressure: $(Bu_3O)_2PO$ 27.79, $(MeO)_2P$ 20.52, $(EtO)_2P$ 24.20, $(PrO)_2P$ 25.91, $(iso-BuO)_2P$ 23.70, $(PhO)_2P$ 42.40, $(BzO)_2PS$ 29.65, $PhPO(OEt)_2$ 31.95, $MePO(OEt)_2$ 30.70, $EtPO(OEt)_2$ 28.65, $FrPO(OEt)_2$ 28.08, $BuPO(OBu)_2$ 27.85, $(MeOCH_2CH_2O)_2P$ 34.79. (2) From γ and the d_s , the exptl. parachors P were detd. and compared with the P calcd. by the procedure of Gililing (C.A. 35, 9493^a; 37, 1308^b; 39, 235, 3986^c) involving "standard value" (SV) for groups and "expansion corrections" (EC) depending on the chain length. Using G.'s $SV = 119.9$ for $((ClO)_2PO$ and 149.8 for $((ClO)_2PS$), the calcd. P agree with the exptl. P for $(PhO)_2PO$ (88.4) and for $(EtO)_2PS$ (43.12) if corrections are applied to allow for the bending of the chains resulting in parallel alignment beyond the β -C atoms and in virtual 5-membered rings O: P-O-C-C-O. Similar agreement is found for $(iso-PrO)_2PO$ and $(iso-BuO)_2PO$. (3) From the exptl. $P = 267.5$ of $(MeO)_2P$, subtraction of 3×55.2 (for 3 Me) and of $EC = 0.66$, leaves for $((ClO)_2P$, $SV = 101.27$; with this value, agreement between the calcd. and the exptl. P for phosphites is obtained if the same corrections are applied as in the case of phosphates. While in the

latter instance the parallel bending of the chains at the β -C atoms could easily be ascribed to polarization of the chains owing to resonance of the P:O bond, in phosphites the same effect can only arise owing to the presence of the electro pair at P; this necessitates assumption of a tetrahedral disposition of valencies around the trivalent P, with the electron pair occupying the 4th corner of the tetrahedron, and leads to postulation of the existence of optically active antipodes of trivalent N, P, and As.


(4) In the case of phosphates of branched alkyls, the uncertainty as to whether the β -C correction should be applied to only one β -C in each chain or to all, was settled in favor of the latter alternative; e.g., for (iso-Pr₂O)PO, the exptl. and calcd. P agree only if the correction is made for all six β -C atoms. The available data do not warrant a decision as to whether the correction for parallelism should be applied only to the principal chain or to both principal and lateral chains. For (MeOCH₂CH₂O)P it is necessary to correct only for the parallelism of one atom (C or O).

(5) From the P of phosphonates, applying the same correction for parallelism as in the case of phosphates, one finds for the group $((\text{C}_2\text{O})_2)(\text{C}_2\text{O})\text{PO}$, $SV = 93.7$. The corrections allowing for the interaction between the resonating P:O bond and the alkyl linked directly to P, are different from those pertaining to the ester alkyls, and, for reasons of analogy, are closer to the corrections applied by Gibling to the bending of chains in esters of carboxylic acids. The structure of phosphonic acid esters resembles closely that of phosphates; it shows 3 virtual δ -membered

APPROVED FOR RELEASE: 06/05/2000

CIA-RDP86-00513R000101920003-5"

ring O.P.C.C.C. (6) Whereas the discrepancy between the exptl. P and the P calcd. according to Gibling is mostly 0.2-0.4% and only exceptionally as high as 0.7%, agreement with the P calcd. according to Samuel (C.A. 38, 35211) is much less satisfactory; in particular, in the case of alkanephosphonic acid esters, the deviation attains 1.5%. Calcd. according to Mumford and Phillips (C.A. 22, 1325; 24, 53) gives errors up to 3.5%. N. Thon



		1ST AND 2ND CODES																3RD AND 4TH CODES															
		PROCESSES AND PROPERTIES INDEX																															
		CA																10															
		<p>Action of 1-bromo-1-nitro hydrocarbons on triethyl phosphate. A. E. Arbusov, B. A. Arbuзов, and B. P. Lugovkin. Bull. acad. sci. U.R.S.S. Classe sci. chim. 1947, 835-8 (in Russian).—RCHE: NO₂ react with (EtO)₃P not along the lines of expected formation of RCH(ONO)₂(PO(OEt))₃, but anomalously with formation of Et₂PO₃ as the only identifiable product, evidently by a mechanism of the type presented by Allen and Wilson (C.A. 34, 2265) for the thermal decompn. of bromonitro derivs. through a biradical mechanism. The over-all reaction may be given by: CH₃BrNO₂ → CH₃⁺ + Br + NO₂ + P(OEt)₃ → CH₃CH₂ + Et₂PO₃ + NO₂ + Et₂PO(OEt)₂. The nitro group does not in itself utilize (EtO)₃P. To 11.1 g. (EtO)₃P in 20 cc. Et₂O was slowly added 8.8 g. CH₃BrNO₂ in 20 cc. Et₂O over 20 min.; the temp. rose spontaneously to 40° and, after the Et₂O was distd., the residue was heated to 100°, when a spontaneous reaction raised the temp. to 100° with evolution of 49.2% EtBr; distn. of the residue gave 65% Et₂PO₃ and 8 g. unidentified solid residue; the off-gas produced in the reaction was unsatd. and presumed to be CO₂. Similar treatment of Et₂PO₃ with an equimol. amt. of MeC₁₁H₂₃NH₃ gave 47% Et₂PO₃. Heating Et₂PO₃ with MeNO₂ gave no reaction. To 28.4 g. Et₂PO₃ at 4° was added slowly 22 g. PhC₆H₅NO₂; spontaneous reaction raised the temp. to 92° and EtBr began to distil; the mass was heated to 180°, when EtBr distn. ceased for a total yield of 74%. Distn. gave 15.6 g. crude Et₂PO₃ and an unidentified product (2.6 g.), b.p. 78-81°, n_D²⁰ 1.5049, which on standing deposited crystals, m. 237-9° (from EtOAc), free from P and Br and analyzed as C₁₁H₂₃N₂O.</p> <p style="text-align: right;">G. M. K.</p>																															
		ASH-SLA METALLURGICAL LITERATURE CLASSIFICATION																E 200013-A-000000															
		120011 120012 120013 120014 120015 120016 120017 120018 120019 120020 120021 120022 120023 120024 120025 120026 120027 120028 120029 120030 120031 120032 120033 120034 120035 120036 120037 120038 120039 120040 120041 120042 120043 120044 120045 120046 120047 120048 120049 120050 120051 120052 120053 120054 120055 120056 120057 120058 120059 120060 120061 120062 120063 120064 120065 120066 120067 120068 120069 120070 120071 120072 120073 120074 120075 120076 120077 120078 120079 120080 120081 120082 120083 120084 120085 120086 120087 120088 120089 120090 120091 120092 120093 120094 120095 120096 120097 120098 120099 1200100 1200101 1200102 1200103 1200104 1200105 1200106 1200107 1200108 1200109 1200110 1200111 1200112 1200113 1200114 1200115 1200116 1200117 1200118 1200119 1200120 1200121 1200122 1200123 1200124 1200125 1200126 1200127 1200128 1200129 1200130 1200131 1200132 1200133 1200134 1200135 1200136 1200137 1200138 1200139 1200140 1200141 1200142 1200143 1200144 1200145 1200146 1200147 1200148 1200149 1200150 1200151 1200152 1200153 1200154 1200155 1200156 1200157 1200158 1200159 1200160 1200161 1200162 1200163 1200164 1200165 1200166 1200167 1200168 1200169 1200170 1200171 1200172 1200173 1200174 1200175 1200176 1200177 1200178 1200179 1200180 1200181 1200182 1200183 1200184 1200185 1200186 1200187 1200188 1200189 1200190 1200191 1200192 1200193 1200194 1200195 1200196 1200197 1200198 1200199 1200200 1200201 1200202 1200203 1200204 1200205 1200206 1200207 1200208 1200209 1200210 1200211 1200212 1200213 1200214 1200215 1200216 1200217 1200218 1200219 1200220 1200221 1200222 1200223 1200224 1200225 1200226 1200227 1200228 1200229 1200230 1200231 1200232 1200233 1200234 1200235 1200236 1200237 1200238 1200239 1200240 1200241 1200242 1200243 1200244 1200245 1200246 1200247 1200248 1200249 1200250 1200251 1200252 1200253 1200254 1200255 1200256 1200257 1200258 1200259 1200260 1200261 1200262 1200263 1200264 1200265 1200266 1200267 1200268 1200269 1200270 1200271 1200272 1200273 1200274 1200275 1200276 1200277 1200278 1200279 1200280 1200281 1200282 1200283 1200284 1200285 1200286 1200287 1200288 1200289 1200290 1200291 1200292 1200293 1200294 1200295 1200296 1200297 1200298 1200299 1200300 1200301 1200302 1200303 1200304 1200305 1200306 1200307 1200308 1200309 1200310 1200311 1200312 1200313 1200314 1200315 1200316 1200317 1200318 1200319 1200320 1200321 1200322 1200323 1200324 1200325 1200326 1200327 1200328 1200329 1200330 1200331 1200332 1200333 1200334 1200335 1200336 1200337 1200338 1200339 1200340 1200341 1200342 1200343 1200344 1200345 1200346 1200347 1200348 1200349 1200350 1200351 1200352 1200353 1200354 1200355 1200356 1200357 1200358 1200359 1200360 1200361 1200362 1200363 1200364 1200365 1200366 1200367 1200368 1200369 1200370 1200371 1200372 1200373 1200374 1200375 1200376 1200377 1200378 1200379 1200380 1200381 1200382 1200383 1200384 1200385 1200386 1200387 1200388 1200389 1200390 1200391 1200392 1200393 1200394 1200395 1200396 1200397 1200398 1200399 1200400 1200401 1200402 1200403 1200404 1200405 1200406 1200407 1200408 1200409 1200410 1200411 1200412 1200413 1200414 1200415 1200416 1200417 1200418 1200419 1200420 1200421 1200422 1200423 1200424 1200425 1200426 1200427 1200428 1200429 1200430 1200431 1200432 1200433 1200434 1200435 1200436 1200437 1200438 1200439 1200440 1200441 1200442 1200443 1200444 1200445 1200446 1200447 1200448 1200449 1200450 1200451 1200452 1200453 1200454 1200455 1200456 1200457 1200458 1200459 1200460 1200461 1200462 1200463 1200464 1200465 1200466 1200467 1200468 1200469 1200470 1200471 1200472 1200473 1200474 1200475 1200476 1200477 1200478 1200479 1200480 1200481 1200482 1200483 1200484 1200485 1200486 1200487 1200488 1200489 1200490 1200491 1200492 1200493 1200494 1200495 1200496 1200497 1200498 1200499 1200500 1200501 1200502 1200503 1200504 1200505 1200506 1200507 1200508 1200509 1200510 1200511 1200512 1200513 1200514 1200515 1200516 1200517 1200518 1200519 1200520 1200521 1200522 1200523 1200524 1200525 1200526 1200527 1200528 1200529 1200530 1200531 1200532 1200533 1200534 1200535 1200536 1200537 1200538 1200539 1200540 1200541 1200542 1200543 1200544 1200545 1200546 1200547 1200548 1200549 1200550 1200551 1200552 1200553 1200554 1200555 1200556 1200557 1200558 1200559 1200560 1200561 1200562 1200563 1200564 1200565 1200566 1200567 1200568 1200569 1200570 1200571 1200572 1200573 1200574 1200575 1200576 1200577 1200578 1200579 1200580 1200581 1200582 1200583 1200584 1200585 1200586 1200587 1200588 1200589 1200590 1200591 1200592 1200593 1200594 1200595 1200596 1200597 1200598 1200599 1200600 1200601 1200602 1200603 1200604 1200605 1200606 1200607 1200608 1200609 1200610 1200611 1200612 1200613 1200614 1200615 1200616 1200617 1200618 1200619 1200620 1200621 1200622 1200623 1200624 1200625 1200626 1200627 1200628 1200629 1200630 1200631 1200632 1200633 1200634 1200635 1200636 1200637 1200638 1200639 1200640 1200641 1200642 1200643 1200644 1200645 1200646 1200647 1200648 1200649 1200650 1200651 1200652 1200653 1200654 1200655 1200656 1200657 1200658 1200659 1200660 1200661 1200662 1200663 1200664 1200665 1200666 1200667 1200668 1200669 1200670 1200671 1200672 1200673 1200674 1200675 1200676 1200677 1200678 1200679 1200680 1200681 1200682 1200683 1200684 1200685 1200686 1200687 1200688 1200689 1200690 1200691 1200692 1200693 1200694 1200695 1200696 1200697 1200698 1200699 1200700 1200701 1200702 1200703 1200704 1200705 1200706 1200707 1200708 1200709 1200710 1200711 1200712 1200713 1200714 1200715 1200716 1200717 1200718 1200719 1200720 1200721 1200722 1200723 1200724 1200725 1200726 1200727 1200728 1200729 1200730 1200731 1200732 1200733 1200734 1200735 1200736 1200737 1200738 1200739 1200740 1200741 1200742 1200743 1200744 1200745 1200746 1200747 1200748 1200749 1200750 1200751 1200752 1200753 1200754 1200755 1200756 1200757 1200758 1200759 1200760 1200761 1200762 1200763 1200764 1200765 1200766 1200767 1200768 1200769 1200770 1200771 1200772 1200773 1200774 1200775 1200776 1200777 1200778 1200779 1200780 1200781 1200782 1200783 1200784 1200785 1200786 1200787 1200788 1200789 1200790 1200791 1200792 1200793 1200794 1200795 1200796 1200797 1200798 1200799 1200800 1200801 1200802 1200803 1200804 1200805 1200806 1200807 1200808 1200809 1200810 1200811 1200812 1200813 1200814 1200815 1200816 1200817 1200818 1200819 1200820 1200821 1200822 1200823 1200824 1200825 1200826 1200827 1200828 1200829 1200830 1200831 1200832 1200833 1200834 1200835 1200836 1200837 1200838 1200839 1200840 1200841 1200842 1200843 1200844 1200845 1200846 1200847 1200848 1200849 1200850 1200851 1200852 1200853 1200854 1200855 1200856 1200857 1200858 1200859 1200860 1200861 1200862 1200863 1200864 1200865 1200866 1200867 1200868 1200869 1200870 1200871 1200872 1200873 1200874 1200875 1200876 1200877 1200878 1200879 1200880 1200881 1200882 1200883 1200884 1200885 1200886 1200887 1200888 1200889 1200890 1200891 1200892 1200893 1200894 1200895 1200896 1200897 1200898 1200899 1200900 1200901 1200902 1200903 1200904 1200905 1200906 1200907 1200908 1200909 1200910 1200911 1200912 1200913 1200914 1200915 1200916 1200917 1200918 1200919 1200920 1200921 1200922 1200923 1200924 1200925 1200926 1200927 1200928 1200929 1200930 1200931 1200932 1200933 1200934 1200935 1200936 1200937 1200938 1200939 1200940 1200941 1200942 1200943 1200944 1200945 1200946 1200947 1200948 1200949 1200950 1200951 1200952 1200953 1200954 1200955 1200956 1200957 1200958 1200959 1200960 1200961 1200962 1200963 1200964 1200965 1200966 1200967 1200968 1200969 1200970 1200971 1200972 1200973 1200974 1200975 1200976 1200977 1200978 1200979 1200980 1200981 1200982 1200983 1200984 1200985 1200986 1200987 1200988 1200989 1200990 1200991 1200992 1200993 1200994 1200995 1200996 1200997 1200998 1200999 12001000 12001001 12001002 12001003 12001004 12001005 12001006 12001007 12001008 12001009 120010010 120010011 120010012 120010013 120010014 120010015 120010016 120010017 120010018 120010019 120010020 120010021 120010022 120010023 120010024 120010025 120010026 120010027 120010028 120010029 120010030 120010031 120010032 120010033 120010034 120010035 120010036 120010037 120010038 120010039 120010040 120010041 120010042 120010043 120010044 120010045 120010046 120010047 120010048 120010049 120010050 120010051 120010052 120010053 120010054 120010055 120010056 120010057 120010058 120010059 120010060 120010061 120010062 120010063 120010064 120010065 120010066 120010067 120010068 120010069 120010070 120010071 120010072 120010073 120010074 120010075 120010076 120010077 120010078 120010079 120010080 120010081 120010082 120010083 120010084 120010085 120010086 120010087 120010088 120010089 120010090 120010091 120010092 120010093 120010094 120010095 120010096 120010097 120010098 120010099 1200100100 1200100101 1200100102 1200100103 1200100104 1200100105 1200100106 1200100107 1200100108 1200100109 1200100110 1200100111 1200100112 1200100113 1200100114 1200100115 1200100116 1200100117 1200100118 1200100119 1200100120 1200100121 1200100122 1200100123 1200100124 1200100125 1200100126 1200100127 1200100128 1200100129 1200100130 1200100131 1200100132 1200100133 1200100134 1200100135 1200100136 1200100137 1200100138 1200100139 1200100140 1200100141 1200100142 1200100143 1200100144 1200100145 1200100146 1200100147 1200100148 1200100149 1200100150 1200100151 1200100152 1200100153 1200100154 1200100155 1200100156 1200100157 1200100158 1200100159 1200100160 1200100161 1200100162 1200100163 1200100164 1200100165 1200100166 1200100167 1200100168 1200100169 1200100170 1200100171 1200100172 1200100173 1200100174 1200100175 1200100176 1200100177 1200100178 1200100179 1200100180 1200100181 1200100182 1200100183 1200100184 1200100185 1200100186 1200100187 1200100188 1200100189 1200100190 1200100191 1200100192 1200100193 1200100194 1200100195 1200100196 1200100197 1200100198 1200100199 1200100200 1200100201 1200100202 1200100203 1200100204 1200100205 1200100206 1200100207 1200100208 1200100209 1200100210 1200100211 1200100212 1200100213 1200100214 1200100215 1200100216 1200100217 1200100218 1200100219 12001002																															

CARBON ELEMENTS		PROCESSES AND PROPERTIES INDEX																																																																				
1ST AND 2ND ORDERS																																																																						
<i>CA</i>																																																																						
<p>Parachors and the structure of dialkyl phosphorous acids N. A. Arbusov and V. S. Vinogradova, <i>Bull. acad. sci. U.R.S.S., Classe sci. chim.</i>, 1947, 617-22 (in Russian).—In continuation of parachor studies on $(RO)_2PO$ and $RPO(OH)_2$ (cf. C.A. 42, 3126), the work was extended to $(RO)_3POH$, with the hope of elucidating the possible tautomeric structures. Using Gibling's method (C.A. 39, 2389), with the group value of PO_2OH being 116.45 (calcd. from P of $(MeO)_2POH$), satisfactory agreement was obtained between observed and calcd. values up to the tBu ester, beyond which a const. deviation of 1.1% was observed. Application of a 2.2 unit correction for each C beyond the β-position gives somewhat better agreement, but such doubling of Gibling's standard β-C correction has no theoretical basis, as long as a monomeric structure is assumed. If a dimeric H-bonded ring structure is assumed, as discussed by Arbusov, et al. (C.A. 41, 4222), the chain interaction correction is applicable to 4 ester chains, i.e., a correction of 8.8 units per every 4 C atoms beyond β-C; i.e., 2.2 units per C atom is explicable. Use of this concept gives the ring-dimer group value for</p>																																																																						
<p>ASB-SEA METALLURGICAL LITERATURE CLASSIFICATION</p>																																																																						
<table border="1"> <thead> <tr> <th colspan="2">TONE STANDARD</th> <th colspan="10">STRONG REL. ONLY ONE</th> </tr> <tr> <th>SEARCHED</th> <th>INDEXED</th> <th colspan="10"></th> </tr> </thead> <tbody> <tr> <td>D</td> <td>S</td> <td>S</td> <td>A</td> <td>V</td> <td>H</td> <td>O</td> <td>I</td> <td>T</td> <td>N</td> <td>M</td> <td>E</td> <td>Y</td> <td>Z</td> <td>W</td> <td>D</td> <td>G</td> </tr> <tr> <td></td> </tr> </tbody> </table>												TONE STANDARD		STRONG REL. ONLY ONE										SEARCHED	INDEXED											D	S	S	A	V	H	O	I	T	N	M	E	Y	Z	W	D	G																		
TONE STANDARD		STRONG REL. ONLY ONE																																																																				
SEARCHED	INDEXED																																																																					
D	S	S	A	V	H	O	I	T	N	M	E	Y	Z	W	D	G																																																						
<p style="text-align: right;">G. M. Kowdappa</p>																																																																						

ARBZOV, B. A.

USSR/Chemistry - Phosphoric Acid
Chemistry - Ethers, Ethyl

Sep/Oct 1947

"The Action of α -Bromonitrohydrocarbons on the Ethyl Ether of Iphosphorous Acid,"
A. E. Arbuzov, B. A. Arbuzov; B. P. Lugovkin, Chem Sci Res Inst immei A. M. Butlerov,
Kazan State U, 3 PP

"Izv Akad Nauk SSSR, Otd Khim Nauk" No 5

Contains experimental data and results, and shows that in each case where bromoni-
tromethane, α -bromonitroethane and α , α -dibromosphenylnitromethane acts on
the ethyl ether of phosphorous acid, the result is triethylphosphate.

IA 53T2

ARBUZOV, B. A.

USSR/Chemistry - Allyl Groups
Chemistry - Esters

Sep/Oct 1947

"Allyl Regroupings," A. N. Fudovik, B. A. Arbuzov, Kazan' State U, 8 pp

"Izv Akad Nauk SSSR, Otd Khim Nauk" No 5

Describes action of sodium and magnesium malonic esters on isomeric methoxy-chloropentanes, and shows that reaction of esters with a secondary chloride gives same products as with a primary chloride.

PA 53T11

ARBUZOV, B. A.

IA 53T7

SSSR/Chemistry - Parachors
Chemistry - Esters

Sep/Oct 1947

"Parachors and the Structure of Phosphoric Acid, Phosphorous Acid and Phosphinous Acid Esters," B. A. Arbu-zov, V. S. Vinogradova, Sci Res Inst imeni A. M. Butlerov, Kazan State U, 13 pp

"Izv Akad Nauk SSSR, Otd Khim Nauk" No 5

Includes data on measurements of parachors of 13 esters of phosphoric, phosphorous and phosphinous acids, and compares them with parachors calculated by Mumford and Phillips and Samuel methods of group values.

53T7

No. 12
1st printing copies
Organophosphorus-tin compounds. I. Synthesis of compounds R_2SnPO_2R' . B. A. Arbusov and A. N. Pudovik (Chem. Inst. Acad. Sci., Kazan). J. Gen. Chem. (U.S.S.R.) 17, 2168-63 (1947).—The Arbusov reaction was used to prep. compds. of the type R_2SnPO_2R' by the interaction of R_2SnX with $P(OR')_3$. The compds. suffer rapid hydrolysis on heating or standing with 5-20% HCl, with rupture of the P-Sn bond; similar cleavage occurs with alk. solns. or with halogens, as well as with AcCl. When 16 g. Me_2SnI and 7 g. $(EtO)_3P$ were heated to 105° a vigorous reaction took place and MeI distd. from the mixt.; the collected distillate was returned and the distn. was repeated at a max. temp. of 110°; 6.2 g. MeI was collected (7.7 g. theoretical); the residue, remaining in the flask, crystd. on standing and represented a 95% yield of $Me_2SnPO(OEt)_2$, m. 91° (from Et_2O), sol. in Et_2O , $CHCl_3$, $MeCO$, $BuOH$, and H_2O . This, on heating 3 hrs. with concd. HCl to 100°, gave Me_2SnCl_2 , m. 107°; standing at room temp. in 5% HCl gave Me_2SnCl_2 , m. 42°, b.p. 45-7°; boiling 3 hrs. with distd. H_2O did not change the product, however; treatment with Cl in $CHCl_3$ soln. for 24 hrs. also gave Me_2SnCl_2 as did similar treatment with AcCl; boiling with 10% aq. KOH gave Me_2SnOH , m. 118°. Heating 23 g. Et_2SnI and 12 g. $(EtO)_3P$ 15 min. to 150° gave 9 g. crude $Et_2SnPO(OEt)_2$, which on distn. gave 3 g. pure product, viscous liquid, b.p. 210-20°, n_D²⁵ 1.4858; this heated with 20% HCl 10 hrs. at 100-70° in a sealed tube gave Et_2SnCl_2 , m. 84.5-85°, while treatment with AcCl and 2 days' standing gave Et_2SnCl_2 , b.p. 98-100°, and a small amt. of unidentified product, b.p. 142-5°, which reduces Fehling soln. $(EtO)_3PONa$ from 4 g. $(EtO)_3POH$ in 30 ml. Et_2O was treated with 8 g. Me_2SnI , boiled 1 hr., the heavy white ppt. sepd., and the Et_2O evapd., to give a yellow oil, which on standing 24 hrs. deposited 0.88 g. $(Me_2SnOH)_2$, Me_2SnI , decomp. 145-65°, which apparently

was formed from the free radical Me_2Sn , produced in the initial reaction with the sodium phosphite. The $(EtO)_3PONa$ from 4.2 g. $(EtO)_3POH$ in abs. Et_2O was treated with 10 g. Et_2SnI (no NaI pptn. was observed), the Et_2O was distd. and was replaced by MeI^{1/2}, the mixt. boiled, and the soln. decanted from the white solid ppt. and distd. to give a colorless product, b.p. 83-4°, n_D²⁵ 1.4842, d₄²⁰ 1.3042, which appeared to be Et_2SnOEt . Heating Ph_2SnBr with $(EtO)_3P$ 0.6 hr. to 170° gave only Ph_2Sn , m. 222°; similarly, when $(EtO)_3PONa$ (from 0.20 g. Na) in Et_2O was mixed with 4.8 g. Ph_2SnBr in benzene, heated 2 hrs., freed of Et_2O , and heated 4 hrs. on a steam bath, the filtered soln. on concn. had an odor of $(EtO)_3P$; addn. of Et_2O pptd. 3.7 g. Ph_2Sn , m. 222°. Heating Ph_2PbBr with $(EtO)_3P$ gave only the disproportionation product, Ph_2Pb , m. 224°. Heating 8.5 g. Et_2PbBr and 4.5 g. $(EtO)_3P$ 6 hrs. at 100° gave some $(EtO)_3P$ and a white amorphous solid, insol. in org. solvents. A similar product is obtained on heating a PhMe soln. of Et_2PbBr , apparently a disproportionation product of Et_2PbBr . Thus, the org.-Pr compds. were incapable of the normal Arbusov reaction. II. Synthesis of compounds of the type $R_2Sn(PO_2R')_2$. B. A. Arbusov and N. P. Grechkin. Ibid. 2166-77.—The Arbusov reaction with R_2SnX_2 gave the expected phosphono-tin compds. of type $R_2Sn(PO_2R')_2$, solids or semisolids, which in most cases could be crystd. from org. solvents. They were sol. in hot $BuOH$ and $CHCl_3$, almost insol. in other solvents, and their crystn. led to severe losses. The high m.p. and poor solv. of the products is explained by their apparent dimeric structure, as shown by Rast mol. wt. detns. of the Et and the Pr compds. The P-Sn bond is readily cleaved on treatment with dil. HCl even at room temp., giving Et_2SnCl_2 , while HBr gives Et_2SnBr ; 10% NaOH gives Et_2SnO , while Cl or Br in $CHCl_3$ soln. give Et_2SnX_2 . The study was extended briefly to reactions of $RSnX_2$ and SnX_4 , which will be reported later. However, the reaction of Me_2SnI with $2(EtO)_3P$ gave, on heating to 85-90°, followed by cooling,

10

an abundant crystn. of a golden yellow solid, identified as an intermediate adduct, $\text{Me}_3\text{Sn}[\text{P}(\text{OEt})_3]_2$, which was an addnl. confirmation of the addn. mechanism of the Arbuzov reaction; this adduct m. 101-3° (evolution of EtI). When 20.2 g. Me_3SnI , and 12.4 g. $(\text{MeO})_2\text{P}$ were heated to 90-100°, a vigorous reaction took place in which 12.5 g. MeI distd. from the mixt.; the glassy residue, dissolved in hot Bu_3OH , gave on cooling 40.6% $\text{Me}_3\text{Sn}(\text{PO}_2\text{Et})_3$, m. 245-7°. An analogous procedure was used to prep. the following: $\text{Et}_2\text{Sn}(\text{PO}_2\text{Et})_3$, m. 249-51°, 40%; $\text{Pr}_2\text{Sn}(\text{PO}_2\text{Pr})_3$, m. 251-3°, 12%; $\text{Et}_3\text{Sn}(\text{PO}_2\text{Et})_3$, m. 202-4°, 10%; $\text{Et}_2\text{Sn}(\text{PO}_2\text{Me})_3$, m. 203.5-05°, 43%; $\text{Me}_3\text{Sn}(\text{PO}_2\text{Et})_3$, glassy solid, no m.p., or yield cited. When $\text{Me}_3\text{Sn}(\text{PO}_2\text{Et})_3$ was slowly added to 6.2 g. Me_3SnI , a brisk reaction occurred; the mixt. was heated 10 min. to 60-80°, and the MeI removed *in vacuo* (the wt. loss was equal to theoretical); the residual crumbly white solid was free of halogen, but due to insol. in org. solvents could not be purified; its behavior with 30% HBr (formation of Me_3SnBr in 50% yield) suggested that it was $\text{Me}_3\text{Sn}(\text{PO}_2\text{Et})_3$. Addn. of 4.9 g. $(\text{EtO})_2\text{P}$ to 5.5 g. SnI_3 , followed by heating to 140° until soln. took place, and removal of EtI *in vacuo* (the wt. loss was equal to theoretical) gave a viscous yellow mass, which could not be purified by crystn. and whose formation of SnCl_3 with hot HCl suggested its

structure as $\text{Sn}(\text{PO}_2\text{Et})_3$, although satisfactory analyses were impossible because of extremely difficult combustion. An Et_2O soln. of 10 g. $(\text{EtO})_2\text{PONa}$ (10 g.), boiled 10 min. with 21.5 g. Et_2SnI_3 in Et_2O , the NaI ppt. filtered off (82.7%), and the filtrate concd., gave 23 g. yellowish liquid which smelled like $(\text{EtO})_2\text{P}$; on standing 3-4 hrs. it deposited 7.7 g. of a white amorphous solid identified as Et_2SnO (80%). The washing of the solid (Bu_3OH , EtOH , and Et_2O) on concn. gave a sirup which on long standing deposited 0.8 g. colorless prisms, m. 173-4°, identified as $\text{Et}_2\text{Sn}(\text{PO}_2\text{Et})_3$, by analysis and the formation of Et_2SnBr_3 with 30% HBr. This product was monomer, as shown by mol. wt. detn. in CHCl_3 . Equimol. amts. of $(\text{MeO})_2\text{PONa}$ and Me_3SnI , gave Me_3SnO as the only identified product (yields Me_3SnCl_3 , m. 102-3°, on treatment with HCl). Similarly, $(\text{EtO})_2\text{PONa}$ and Et_2SnCl_3 in Et_2O gave 50% Et_2SnO . The oxide is probably the result of atm. O oxidation of the free radical R_3Sn , which is formed in the primary action. Na (1.1 g.) and 6.6 g. $(\text{EtO})_2\text{PONa}$ in ether with 8.3 g. Me_3SnI , immediately formed a ppt. of NaI and when boiled 0.25 hr. gave 97.5% NaI; evapn. of the filtrate gave a mobile liquid yielding, on distn., 1.8 g. mixed $(\text{EtO})_2\text{P}$ and $(\text{EtO})_2\text{POH}$, after the removal of which the residue decompd. on attempted distn. *in vacuo*; the solid residue was insol. in org. solvents or in HCl (the reaction is still under study at this time).

G. M. Kosolapoff

TRANSLATION: 446940

"APPROVED FOR RELEASE: 06/05/2000

CIA-RDP86-00513R000101920003-5

ARBUZOV, B. A. and N. P. Grechkin

"Tin-Phosphorus Organic Compounds and Synthesis of Compounds to Type R₂Sn."
Zhurnal Obshchei Khimii, Vol. XVII, No. 12, 1947, pp. 2166-77.

TRANSLATION Available (456199)

APPROVED FOR RELEASE: 06/05/2000

CIA-RDP86-00513R000101920003-5"

ABUZOV, B. A. (and Pudovik, A. N. (?)

"Obtaining Several Esters of Triarylethylphosphinic Acid O Poluchanii Nekotorikhovoi Kisioty,"(identified with Pudovik, A. N.) Journ. of Gen. Chem., XVII, 1947, 2193-2148, 2158-2165.

ARBUZOV, B. A.

PA 52T15

USER/Chemistry - Boric Acid
Chemistry - Esters

Oct 1947

"Parachors of Substituted Esters of Boric Acid,"
B. A. Arbuzov, V. S. Vinogradova, Sci Res Inst imeni
A. M. Butlerov, Kazan State U imeni V. I. Lenin, 4 pp

"Dok Akad Nauk SSSR" Vol LVIII, No 1

Concludes from data that parachors show the structure
of esters of boric acid, which is in complete
agreement with data of electro organic research.
Favorable concurrence found between calculated and
actual significance of parachors.

52T15

ARBUZOV, B. A.

PA 58T11

USSR/Chemistry - Phosphoric Acid, Dialkyl Jan 1947
Chemistry - Parachors

"Parachor and Structure of Dialkylphosphoric Acid,"
B. A. Arbuzov, V. S. Vinogradova, Sci Res Chem Inst
imeni A. M. Butlerov, Kazan State U, 2 pp

"Dok Akad Nauk SSSR, Nova Ser" Vol LV, No 1

Describes experiments which produce data on para-
chors of dialkylphosphoric acid, fully confirming
results on polymerizing ability of this acid in
liquid condition.

58T11